

Vecuronium

Newborn use only

2025

Alert	High-alert medication: High risk of causing significant patient harm when used in error. ¹ This drug should be administered in the presence of personnel trained in advanced airway management. Reversal agents should be immediately available (see Special Comments). Suggest regular cessation of infusion for a few to several hours, possibly every 24 hours (commonly referred to as a 'drug holiday') to assess the need for continued paralysis and adequacy of sedation or analgesia. Following cessation, the line should be adequately flushed to avoid unintended paralysis during later use of the line. Eye lubricant should be used whilst patient is receiving vecuronium.					
Indication	1. Skeletal muscle relaxation or paralysis in mechanically ventilated infants. 2. For elective endotracheal intubation.					
Action	Nondepolarizing neuromuscular blockers are competitive acetylcholine (ACh) antagonists that bind directly to nicotinic receptors on the postsynaptic membrane, thus blocking the binding of ACh so the motor endplate cannot depolarize. Onset of action is 1-2 minutes; duration of action is 30-40 minutes.					
Drug Type	Non-depolarising neuromuscular blocking agent.					
Trade Name	Vecure Powder for injection, Vecure Sun Powder for injection					
Presentation	10 mg of vecuronium bromide in glass vial (powder for reconstitution)					
Dose	Intubation IV bolus – 100 microgram/kg Muscle relaxation** Intermittent IV bolus 100 microgram/kg (30-150 microgram/kg) IV push every 1-2 hours as required. Continuous IV infusion (with or without loading dose) 100 microgram/kg/hour (60-200 microgram/kg/hour). Start 20 minutes post bolus recovery. Titrate in 10% dose increments until desired neuromuscular blockade is achieved. * Provide eye protection and instil lubricating eye drops every 2 hours. # Sensation remains intact; additional sedation & analgesia should be used for painful procedures.					
Dose adjustment	Therapeutic hypothermia – No information. ECMO – No information. Renal impairment – No specific dose adjustment. However, duration of action may be prolonged. Hepatic impairment – No specific dose adjustment. However, hepatic impairment decreases clearance resulting in prolonged duration of action.					
Route	IV					
Maximum Dose	IV bolus: 200 microgram/kg IV infusion: 200 microgram/kg/hour.					
Total cumulative dose						
Preparation	IV bolus: Add 5 mL water for injection to 10 mg of vecuronium powder for reconstitution vial to make a 2 mg/mL solution. Further dilute: From this vial, draw up 2 mL (4 mg of vecuronium) and add to 2 mL of sodium chloride 0.9% to make a final volume of 4 mL with a concentration of 1 mg/mL [1000 microgram/mL]. IV infusion: Infant <1.5 Kg <table><tr><th>Infusion rate</th><th>Prescribed amount</th></tr><tr><td>1ml/hour = 400 microgram/kg/hour</td><td>20 mg/kg vecuronium and make up to 50 ml</td></tr></table>		Infusion rate	Prescribed amount	1ml/hour = 400 microgram/kg/hour	20 mg/kg vecuronium and make up to 50 ml
Infusion rate	Prescribed amount					
1ml/hour = 400 microgram/kg/hour	20 mg/kg vecuronium and make up to 50 ml					

	First Step: Add 5 mL water for injection to 10 mg vecuronium powder for reconstitution vial to make a 2 mg/mL solution (Note: May need multiple vials based on bodyweight). Further dilute: Draw up 10 mL/kg of this solution (20 mg/kg) and dilute with glucose 5% or sodium chloride 0.9% to make a final volume of 50mL with a concentration of 0.25 mL/hour = 100 microgram/kg/hour. IV bolus dose from this solution: 0.25 mL = 100 microgram/kg. Infant ≥ 1.5 Kg <table border="1"> <thead> <tr> <th>Infusion rate</th><th>Prescribed amount</th></tr> </thead> <tbody> <tr> <td>1ml/hour = 200 microgram/kg/hour</td><td>10 mg/kg vecuronium and make up to 50 ml</td></tr> </tbody> </table> First Step: Add 5 mL water for injection to 10 mg vecuronium powder for reconstitution vial to make a 2 mg/mL solution (Note: May need multiple vials based on bodyweight). Further dilute: Draw up 5 mL/kg of this solution (10 mg/kg) and dilute with glucose 5% or sodium chloride 0.9% to make a final volume of 50mL with a concentration of 0.5 mL/hour = 100 microgram/kg/hour. IV bolus dose from this solution: 0.5 mL = 100 microgram/kg.	Infusion rate	Prescribed amount	1ml/hour = 200 microgram/kg/hour	10 mg/kg vecuronium and make up to 50 ml
Infusion rate	Prescribed amount				
1ml/hour = 200 microgram/kg/hour	10 mg/kg vecuronium and make up to 50 ml				
Administration	IV bolus: Administer over several seconds. IV infusion via syringe pump. Flush line adequately after each dose with sodium chloride 0.9% to avoid unintended paralysis and/or incompatibility with other medications during later use of the line.				
Monitoring	Continuous cardio-respiratory and pulse oximetry monitoring. Close monitoring of neuromuscular function, sedation, and blood pressure (invasive or non-invasive). Monitor electrolytes and renal function. Monitor injection site for signs of extravasation.				
Contraindications	Hypersensitivity to vecuronium or any component of the formulation. Cross-sensitivity with other neuromuscular-blocking agents may occur; use with extreme caution in patients with previous anaphylactic reactions. Severe electrolyte abnormalities.				
Precautions	Avoid prolonged usage. Factors which can increase duration of neuromuscular blockade: Acidosis, hypothermia, neuromuscular disease, hepatic disease, hypokalaemia, hypermagnesaemia, renal failure, and younger age. Vecuronium is lipid soluble and is predominantly excreted via the liver so poor liver function can cause prolonged effects. Factors which can decrease duration of neuromuscular blockade: Alkalosis and hyperkalaemia. Use cautiously in neonates with hepatic or renal impairment and in neonates with fluid and electrolyte imbalance. Suggest regular cessation of infusion, possibly every 24 hours (commonly referred to as 'drug holiday') to assess the need for continued paralysis and adequacy of sedation or analgesia. Monitoring of fluid balance is essential due to risk of fluid retention. Aminoglycosides & general anaesthetics can increase (potentiate) duration of neuromuscular blockade. Corticosteroids: Concomitant use with corticosteroids has been reported to be associated with development of acute quadriplegic myopathy syndrome (AQMS) in adults.³ However, Recent trials provided no evidence for increased risk of neuromyopathy in patients with sepsis or acute respiratory distress syndrome (ARDS) with the use of corticosteroids or neuromuscular blockers.¹⁷ Adrenaline (epinephrine) can reduce (antagonise) duration of neuromuscular blockade.				

Drug Interactions	Antimicrobials like aminoglycosides, tetracyclines, polymyxins, and clindamycin can potentiate neuromuscular blockade.³ Inhaled anaesthetics can potentiate neuromuscular blockade.³ Anti-epileptics can make patients resistant to vecuronium.³ Local anaesthetics can potentiate neuromuscular blockade.³ Aminoglycosides & general anaesthetics can increase duration of neuromuscular blockade. Corticosteroids: Concomitant use with corticosteroids has been reported to be associated with development of acute quadriplegic myopathy syndrome (AQMS) in adults.³ However, Recent trials provided no evidence for increased risk of neuromyopathy in patients with sepsis or acute respiratory distress syndrome (ARDS) with the use of corticosteroids or neuromuscular blockers.¹⁷ Dexamethasone and hydrocortisone may result in decreased vecuronium effectiveness, prolonged muscle weakness, and myopathy.³ Adrenaline (epinephrine) can reduce (antagonise) duration of neuromuscular blockade.
Adverse Reactions	Hypoxaemia may occur because of inadequate ventilation and deterioration in pulmonary mechanics. Hypotension and bradycardia, particularly when used in combination with opioids. Prolonged paralysis after long-term use. Rare: Anaphylactic reaction and tachycardia.
Overdose	Supportive measure: Ventilatory support and sedation. Reversal of neuromuscular blockade can be achieved by neostigmine (refer to special comments). AUSTRALIA: Contact the Poisons Information Centre on 13 11 26 for information on the management of overdose NEW ZEALAND: Contact the National Poisons Centre on 0800 764 766 for information on the management of overdose.
Compatibility	Fluids: ³ glucose 5%, sodium chloride 0.9%. PN at Y site: Compatible with 2 in 1 solution (Amino acid-glucose-trace element mixture). No information on lipid emulsions. Y-site:³ Adrenaline (epinephrine), alprostadil, amikacin sulfate, aminophylline, amiodarone, ampicillin, atenolol, azithromycin, aztreonam, caffeine citrate, calcium chloride, calcium gluconate, cefazolin, ceftazidime, ceftriaxone, ciprofloxacin, clindamycin, dexamethasone sodium phosphate, dexmedetomidine, digoxin, diltiazem, dobutamine, dopamine, enalaprilat, adrenaline (epinephrine), erythromycin lactobionate, esmolol, fentanyl, fluconazole, fluorouracil, fosphenytoin, gentamicin, glycopyrrolate, heparin, hydralazine, hydrocortisone sodium succinate, insulin (regular), isoprenaline, labetalol, lidocaine, linezolid, lorazepam, magnesium sulfate, meropenem, metoprolol, metronidazole, midazolam, milrinone, morphine, naloxone, nifedipine, nitroglycerin, norepinephrine, octreotide, ondansetron, pamidronate, pentoxifylline, phenobarbital, phenylephrine, potassium acetate, potassium chloride, propofol at vecuronium concentrations of ≤ 1 mg/mL, propranolol hydrochloride, ranitidine hydrochloride, remifentanyl, sodium acetate, sodium bicarbonate, sodium nitroprusside, sodium phosphate, streptozocin, succinylcholine, tacrolimus, theophylline, ticarcillin disodium/clavulante potassium, tigecycline, tobramycin, sulfamethoxazole/trimethoprim, and vancomycin hydrochloride, vasopressin, verapamil, voriconazole, zidovudine, and zoledronic acid.
Incompatibility	Fluids: No information. No information on lipid emulsions. Y site:³ Aciclovir, amphotericin B (all compounds), cefepime, cefotaxime, diazepam, furosemide, ganciclovir, ibuprofen lysine, imipenem/cilastatin sodium, methylprednisolone sodium succinate, micafungin sodium, pantoprazole, phenytoin, piperacillin sodium, piperacillin-tazobactam, propofol at vecuronium concentrations >1 mg/mL, sulbactam/durlobactam, and thiopental sodium.
Stability	Diluted solution stable for up to 24 hours. Discard any unused solution.
Storage	Store below 25°C. Protect from light. Store in accordance with local policies.
Excipients	Citric acid, dibasic sodium phosphate, sodium hydroxide and/or phosphoric acid, and mannitol.
Special Comments	Muscle relaxation is reversed by neostigmine (50 microgram/kg) and atropine (20 microgram/kg). Sugammadex is being increasingly used with extrapolated information from other populations. Sensation remains intact: sedation & analgesia should be used for painful procedures. Provide eye protection and instil lubricating eye drops every 2 hours. Vecuronium produces less tachycardia and hypotension when compared with pancuronium.^{15,16}

	The neuromuscular blockade of vecuronium is of shorter duration than that of pancuronium.^{15,16} Prior administration of suxamethonium shortens onset and may increase depth of blockade; reduce dose and give vecuronium only after recovery from suxamethonium-induced neuromuscular blockade.
Evidence	Background Nondepolarizing neuromuscular blocking agents (NMBA) can be classified into 2 classes: steroidal (rocuronium, vecuronium, pancuronium) or benzylisoquinoline (mivacurium, atracurium, cisatracurium). Nondepolarizing neuromuscular blockers are competitive acetylcholine (ACh) antagonists that bind directly to nicotinic receptors on the postsynaptic membrane, thus blocking the binding of ACh so the motor endplate cannot depolarize. This leads to skeletal muscle paralysis.² Paralysis occurs sequentially because of the differing sensitivity of muscles to NMBAs as well as blood flow to the area. Generally, paralysis begins with smaller, fast twitch muscles such as the eyes and larynx, then affects the limbs, neck, trunk, and upper airway, and eventually progresses to the intercostals and diaphragm until respiration terminates. Recovery from paralysis occurs in the reverse order with function of the diaphragm returning first.⁴ Vecuronium: vecuronium is structurally related to pancuronium. It has a greater potency, shorter duration of action, lack significant cardiovascular effects (tachycardia), and less cumulative properties. Vecuronium produces less tachycardia and hypotension when compared with pancuronium.^{15,16} The neuromuscular blockade of vecuronium is of shorter duration than that of pancuronium.^{15,16} Time of onset of action is 90-120 seconds after IV bolus, with a duration of effect that lasts only 30-40 minutes. Intermittent bolus dosing would need to be so frequent (i.e., every 30 to 60 minutes) that continuous IV infusion is preferred over intermittent boluses to maintain paralysis in ventilated infants.⁵ In comparison, rocuronium is an analogue of vecuronium with a more rapid onset of action (20-100 seconds), but less potent than vecuronium, and hence larger doses (example, 600 microgram/kg of rocuronium, compared to 100 microgram/kg of vecuronium). Tachycardia is more frequent with rocuronium, while vecuronium lacks this effect in regular doses.⁴ The neuromuscular blockade effect of vecuronium is stronger and lasts longer in neonate than infant or adult.⁶ The 95% effective dose (ED₉₅) for NMBDs specifies the dose that produces 95% twitch depression in 50% of individuals. ED₉₅ of vecuronium in neonates (47 microgram/kg) is 40% less than in children (81 microgram/kg) meaning less dose is needed in neonates compared to children.⁷ Efficacy Muscle relaxation Two prospective studies by Meretoja et al in 1988 and 1989 determined the dose responses with vecuronium bolus and continuous infusion in paediatric population.^{7,8} The bolus dose required to achieve effective neuromuscular blockade in neonates was 40% less than in children. The median maintenance dose of 0.1 mg/kg is required in neonates to maintain 1 hour of neuromuscular blockade, in comparison to 0.217 mg/kg/hour in children 3-10 years old. Fitzpatrick et al studied vecuronium to facilitate paralysis in mechanically ventilated paediatric population (4 neonates, and 11 infants and children). A loading dose of 0.1 mg/kg was followed by an infusion of 0.1 mg/kg/hour. The titration rate was adjusted to maintain a neuromuscular block of approximately 90% as assessed by the presence of one response to a train of four (TOF) stimulation. The duration of the infusions varied from 9.5 to 179 hours. Mean recovery times after stopping the infusion were 51.7 (±17.6) and 46.8 (± 16.5) minutes for the children and neonates respectively. No adverse cardiovascular or toxic effects were noted.⁹ Fisher et al determined the recovery period (time from injection to 90% recovery) after a bolus of vecuronium. Recovery was longest for infants (73±27 minutes), compared to children (35±6 minutes).¹⁰ The longer recovery period in neonates is thought to be due higher volume of distribution. Hodges et al evaluated the appropriate vecuronium infusion rates in 12 neonates/infants and 18 children using train of four (TOF) monitoring. Neonates and infants required 45% less vecuronium (mean infusion rate 0.54 mg/kg/hour) than older children (0.99 mg/kg/hour) and had faster recovery (45 min vs 65 min), with no evidence of prolonged weakness.¹¹ Safety

	Adults with hepatic and renal failure have been shown to experience prolonged neuromuscular blockade.^{12,13} Pharmacokinetics Hepatobiliary clearance is the primary route of elimination, accounting for approximately 50% of the dose. Vecuronium is metabolised rapidly in the liver to 3-desacetyl-vecuronium, which is 50–70% as potent as the parent compound. This metabolite is cleared primarily by renal elimination. Approximately 20–30% of vecuronium is excreted unchanged in urine.^{9,11,12} Time of onset of action is 90-120 seconds after IV bolus, with a duration of effect that lasts only 30-40 minutes. (prolonged with higher doses and in preterm infants).^{5,14}
Practice points	Eye lubrication should be applied to all patients.
References	1. Clinical Excellence Commission. Neuromuscular blocking agents. https://www.cec.health.nsw.gov.au/keep-patients-safe/medication-safety/high-risk-medicines/neuromuscular-blocking-agents. Accessed on 7 October 2024. 2. Cook D, Simons DJ. Neuromuscular Blockade. [Updated 2023 Nov 13]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: https://www.ncbi.nlm.nih.gov/books/NBK538301/. 3. MerativeTM Micromedex® Complete IV Compatibility (electronic version). Merative, Ann Arbor, Michigan, USA. Available at: https://www.micromedexsolutions.com/ (cited: Oct/8/2024). 4. Warr J, Thiboutot Z, Rose L, Mehta S, Burry LD. Current Therapeutic Uses, Pharmacology, and Clinical Considerations of Neuromuscular Blocking Agents for Critically Ill Adults. <i>Annals of Pharmacotherapy</i>. 2011;45(9):1116-26. 5. Kandasamy J, Carlo WA. Pharmacologic therapies IV. In <i>Assisted ventilation of the neonate</i>. 2017. Elsevier Inc. 6. Meakin GH. Neuromuscular blocking drugs in infants and children. <i>Continuing Education in Anaesthesia Critical Care & Pain</i>. 2007;7(5):143-7. 7. Meretoja OA, Wirtavuori K, Neuvonen PJ. Age-dependence of the dose-response curve of vecuronium in pediatric patients during balanced anesthesia. <i>Anesthesia & Analgesia</i>. 1988;67(1):21-6. 8. Meretoja O. Is vecuronium a long-acting neuromuscular blocking agent in neonates and infants? <i>British Journal of Anaesthesia</i>. 1989;62(2):184-7. 9. Fitzpatrick KT, Black GW, Crean PM, Mirakhor RK. Continuous vecuronium infusion for prolonged muscle relaxation in children. <i>Can J Anaesth</i>. 1991;38:169-74. 10. Fisher DM, Miller RD. Neuromuscular effects of vecuronium (ORG NC45) in infants and children during N2O, halothane anesthesia. <i>Anesthesiology</i>. 1983;58(6):519-23. 11. Hodges U. Vecuronium infusion requirements in paediatric patients in intensive care units: the use of acceleromyography. <i>British journal of anaesthesia</i>. 1996;76(1):23-8. 12. Bencini A, Scaf A, Sohn Y, Kersten-Kleef U, Agoston S. Hepatobiliary disposition of vecuronium bromide in man. <i>British Journal of Anaesthesia</i>. 1986;58(9):988-95. 13. Lynam DP, Cronnelly R, Castagnoli KP, Canfell PC, Caldwell J, Arden J, et al. The pharmacodynamics and pharmacokinetics of vecuronium in patients anesthetized with isoflurane with normal renal function or with renal failure. <i>Anesthesiology</i>. 1988;69(2):227-31. 14. Johnson PN, Miller J, Gormley AK. Continuous-infusion neuromuscular blocking agents in critically ill neonates and children. <i>Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy</i>. 2011;31(6):609-20. 15. Basta SJ, Savarese JJ, Ali HH et al. Vecuronium does not alter serum histamine within the clinical dose range. <i>Anesthesiology</i> 1983;59:A273. 16. Son SL, Waud BE, Waud DR. A comparison of the neuromuscular blocking and vagolytic effects of ORG NC45 and pancuronium. <i>Anesthesiology</i> 1981;55:12–18. 17. Annane D. What is the evidence for harm of neuromuscular blockade and corticosteroid use in the intensive care unit? <i>Seminars in respiratory and critical care medicine</i> 2016;37:51-56.

VERSION/NUMBER	DATE
Original 1.0	10/04/2017
Version 2.0	27/06/2019
Version 3.0	17/10/2024
Version 3.0 (minor errata)	1/11/2024
Version 4.0	10/07/2025
Current 4.0 (minor errata)	29/09/2025
REVIEW	10/07/2030

Authors Contribution of the current version

Current version authors	Srinivas Bolisetty, Rebecca O'Grady, Nilkant Phad
Evidence Review	Srinivas Bolisetty, Rebecca O'Grady, Nilkant Phad
Expert review	
Nursing Review	Benjamin Emerson-Parker
Pharmacy Review	Rebecca O'Grady
ANMF Group contributors	Bhavesh Mehta, Rebecca Barzegar, Mohammed Irfan Azeem, Thao Tran, Cindy Chen, Susannah Brew, Michelle Jenkins, Celia Cunha Brites, Bryony Malloy, Renae Gengaroli, Samantha Hassall, Amber Seigel, Emma Watson, Tiffany Kwan, Charles Tian
Final editing	Srinivas Bolisetty
Electronic version	Thao Tran, Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty

Citation for the current version

Bolisetty S, O'Grady R, Phad N, Emerson-Parker B, Mehta B, Barzegar R, Azeem MI, Tran T, Chen C, Brew S, Jenkins M, Brites CC, Malloy B, Gengaroli R, Hassall S, Watson E, Kwan T, Seigel A, Callander I. Vecuronium. Consensus formulary by the Australasian Neonatal Medicines Formulary group. Version 4, dated 10 July 2025. www.anmfonline.org