

Alert	Check ampoule carefully as an adult 10 mg ampoule (Konakion MM Adult) is also available. Konakion MM Paediatric is recommended. Vitamin K Deficiency Bleeding is also known as Haemorrhagic Disease of Newborn (HDN).
Indication	Prophylaxis and treatment of vitamin K deficiency bleeding (VKDB) Neonatal cholestasis
Action	Promotes the activation of blood coagulation Factors II, VII, IX, and X in the liver
Drug type	Fat soluble vitamin
Contraindications	Oral prophylaxis is contraindicated in infants who are: preterm, unwell, on antibiotics, have cholestasis, or have diarrhoea. Oral prophylaxis is contraindicated in infants of mothers who are on anticonvulsants including phenytoin, barbiturates, carbamazepine, rifampicin, and vitamin K antagonists including warfarin and phenindione.
Precautions	Efficacy of treatment is decreased in patients with liver disease.
Dose	<p>IM prophylaxis (Recommended route)</p> <ul style="list-style-type: none"> Birthweight ≥ 1500 g: 1 mg (0.1 mL) as a single dose at birth. Birthweight <1500 g: 0.5 mg (0.05 mL) as a single dose at birth. <p>Oral prophylaxis¹ 2 mg (0.2 mL) for 3 doses:</p> <ul style="list-style-type: none"> First dose: at birth. Second dose: day 3 of life (up to 5th days of life). Third dose: during 4th week (day 22-28 of life) if the infant is fully breastfed (if fed mostly by formula a third dose is not required). It is imperative that the third dose is given no later than 4 weeks after birth as the effect of earlier doses decreases after this time. Repeat the oral dose if the infant vomits within an hour of an oral dose or if diarrhoea occurs within 24 hours of administration. Each facility must develop a local strategy to ensure administration of the third dose, as infants are typically in the community by week four and the risk of non-compliance is high. <p>IV Prophylaxis⁵</p> <ul style="list-style-type: none"> May be given in sick infants if unable to give IM or orally. 0.3 mg/kg (0.2-0.4 mg/kg) as a single dose as a slow bolus (maximum 1 mg/minute). Must be administered in the presence of a medical officer. Dose may be repeated weekly. <p>IV treatment of Vitamin K Deficiency Bleeding (VKDB)</p> <ul style="list-style-type: none"> 1 mg IV as a slow bolus. Dose may be repeated in 4–6 hours if required. Must be administered in the presence of a medical officer. <p>Neonatal cholestasis</p> <ul style="list-style-type: none"> Refer to vitamins in cholestasis formulary.
Dose adjustment	Therapeutic hypothermia: No dose adjustment Renal impairment: No dose adjustment Hepatic impairment: No dose adjustment. ECMO: No information.
Maximum dose	No information in neonates. Doses up to 10mg have been reported in paediatrics. Maximum rate (IV): 1 mg/minute
Route	IM, Oral, IV
Trade name	Konakion MM Paediatric
Presentation	2 mg/0.2 mL ampoule
Preparation	IM and Oral: Administer undiluted. IV: Draw up 0.2 mL (2 mg) of Konakion MM Paediatric and add 1.8 mL of glucose 5% or sodium chloride 0.9% to make a 1 mg/mL solution (ANMF consensus).

Compatibility	<p>Fluids⁸: Glucose 5% (use immediately), glucose 10%, sodium chloride 0.9%, sodium chloride 0.45%.</p> <p>Amino acids at Y-site: Variable</p> <p>Lipids at Y-site: no information</p> <p>Y-site⁸: Amikacin, aminophylline, ascorbic acid, atracurium, atropine, azathioprine, aztreonam, benzylpenicillin, calcium chloride, calcium gluconate, cefazolin, cefotaxime, ceftazidime, ceftriaxone, cefuroxime, clindamycin, dexamethasone, dopamine, doxycycline, enalaprilat, adrenaline (epinephrine), epoetin alfa, erythromycin lactobionate, fentanyl, furosemide (frusemide), ganciclovir, gentamicin, heparin sodium, hydrocortisone, indomethacin, insulin regular, isoproterenol, labetalol, lidocaine, midazolam, morphine, naloxone, nitroglycerin, nitroprusside sodium, norepinephrine, oxacillin, penicillin G potassium, penicillin G sodium, phenobarbital (phenobarbitone), piperacillin, potassium chloride, propranolol, protamine, pyridoxine, ranitidine, sodium bicarbonate, streptokinase, succinylcholine, thiamine, ticarcillin, ticarcillin-clavulanate, tobramycin, tolazoline, urokinase, vancomycin, vasopressin, and verapamil.</p> <p>Variable compatibility⁸: Amphotericin B conventional colloidal, ampicillin, dobutamine, hydralazine, and methylprednisolone.</p>
Incompatibility	Y-site ⁸ : Diazepam, diazoxide, magnesium sulfate, phenytoin, and sulfamethoxazole-trimethoprim.
Administration	<p>IM</p> <p>Inject into the thickest part of the vastus lateralis (anterolateral thigh). Oral</p> <p>Injection solution can be administered orally via dispenser provided.</p> <p>Dose is to be repeated if infant spits out or vomits within an hour of an oral dose or if diarrhoea occurs within 24 hours of administration. Check with medical officer for advice.</p> <p>IV</p> <p>Slow bolus. Maximum rate 1 mg/minute. Must be administered in the presence of a medical officer.</p>
Monitoring	Prothrombin time when treating clotting abnormalities (a minimum of 2 to 4 hours is needed for measurable improvement).
Drug interactions	Co-administration of anticonvulsants can impair the action of vitamin K ₁ .
Adverse reactions	<p>Pain, swelling, and erythema at IM injection site.</p> <p>Severe hypersensitivity reactions, including death have been reported with rapid IV administration.</p>
Stability	Use immediately.
Storage	Store below 25°C. Protect from light.
Excipients	Glycocholic acid, lecithin, sodium hydroxide, hydrochloric acid
Special comments	The risk of childhood cancer is not increased by IM administration of vitamin K ₁ .
Evidence	<p>Overview</p> <p>All newborn infants have a relative vitamin K deficiency at birth. Vitamin K₁ crosses the placenta poorly resulting in low foetal plasma concentrations of the vitamin, with a 30:1 maternal-infant gradient. Human breast milk contains relatively low concentrations of vitamin K₁ (1 to 2 mg/L). Relative deficiency of vitamin K₁, particularly in exclusively breastfed infants can lead to Vitamin K Deficiency Bleeding (VKDB), previously known as Haemorrhagic Disease of Newborn (HDN).⁽¹⁾ VKDB is rare but potentially fatal disorder. The overall annual incidence in Australia is 0.84 per 100000 livebirths.¹¹ VKDB is classified into early, classic, and late, based on the age of presentation. Early VKDB, occurring within the first 24 hours of life, is rare and confined to infants born to mothers who have received medications that interfere with vitamin K metabolism. Classic VKDB occurs between 2nd and 7th day of life. Late VKDB occurs from eight days to six months after birth, with most presenting at one to three months of life. Vitamin K prophylaxis at birth is recommended to prevent classic VKDB.^{1-3,9-11}</p> <p>VKDB trends</p> <p>2020 Australian Paediatric surveillance study found the annual incidence of any VKDB was 0.84/100 000 live births and the late VKDB was 0.61/100000 live births. The proportion of parents who refused vitamin K was significantly higher during 2006–2017 compared with the period 1993–2005. In the 5 years 2013–2017, among the 14 cases of VKDB reported in this period, 10 (71%) did not receive vitamin K due to parental refusal.¹¹</p> <p>2026 Swedish population study showed increasing trends of infants not receiving IM vitamin K prophylaxis at birth (from 0.66% in 2006 to 1.5% in 2021). In this study, Infants without IM vitamin K had 1.54-fold higher odds (aOR, 1.52; 95% CI, 1.27-1.81) of bleeding and 3.18-fold higher odds (aOR, 2.91;</p>

	<p>95%CI, 2.13-3.96) of an intracranial bleeding episode during infancy, compared with infants who received intramuscular vitamin K.¹²</p> <p>Efficacy</p> <p>Vitamin K prophylaxis for VKDB in neonates: Cochrane review by Puckett et al. found that a single dose (1 mg) of intramuscular vitamin K₁ after birth is effective in the prevention of classic VKDB. Either intramuscular or oral (1 mg) vitamin K prophylaxis improves biochemical indices of coagulation status at 1–7 days. Neither intramuscular nor oral vitamin K₁ has been tested in randomised trials with respect to effect on late VKDB. When three doses of oral vitamin K₁ are compared to a single dose of IM vitamin K₁, the plasma vitamin K₁ concentrations are higher in the oral group at two weeks and two months, but, again, there is no evidence of a difference in coagulation status.² (LOE II, GOR B)</p> <p>Vitamin K prophylaxis for VKDB in preterm neonates: Cochrane review by Ardell et al. found only one RCT that compared IV to IM administration of vitamin K and compared various dosages of vitamin K. Three different prophylactic regimes of vitamin K (0.5 mg IM, 0.2 mg IM, or 0.2 mg IV) were given to infants less than 32 weeks' gestation. There was no statistically significant difference in vitamin K levels in the 0.2 mg IV group when compared to 0.2 or 0.5 mg IM groups on day 5. By day 25, vitamin K₁ levels had declined in all the groups, but infants who received 0.5 mg IM had higher levels of vitamin K₁ than either the 0.2 mg IV group or the 0.2 mg IM group. Since there is no available evidence that vitamin K is harmful or ineffective and since vitamin K is an inexpensive drug, authors concluded to follow the recommendations of expert bodies and give vitamin K to preterm infants.³</p> <p>Treatment of VKDB: Any infant suspected of VKDB should receive immediate intravenous vitamin K replacement. It is standard practice to administer a dose of 1 mg which will usually result in correction within a few hours. (LOE IV; GOR C) Intravenous vitamin K can be associated with anaphylactoid reactions and should be administered by slow intravenous injection; if venous access cannot be established it can be given subcutaneously, the intramuscular route being avoided in the presence of a coagulopathy.⁴</p> <p>Pharmacokinetics</p> <p>In healthy, fully breast-fed, newborn babies, significantly higher plasma vitamin K₁ concentrations were reported several weeks after IM as compared to oral vitamin K₁. Half-life of oral and intramuscular vitamin K₁ were considerably longer in newborn infants (median 76 hours; range 26 to 193 hours)^{5, 6} compared to adults (6 hours; range 2–26 hours)⁽⁷⁾. Re-dosing of oral vitamin K₁ is recommended by 1 month in breast fed infants.⁶ (LOE II GOR B)</p> <p>In preterm infants and sick infants unable to receive intramuscular vitamin K₁, 0.3 mg/kg intravenously resulted in similar serum concentrations as oral administration of 3 mg vitamin K₁ and intramuscular administration of 1.5 mg vitamin K₁ which then supports recommendation for intravenous 0.4 mg/kg phytomenadione - vitamin K₁ - Konakion MM Paediatric in infants unable to receive oral or intramuscular vitamin K₁.⁵ (LOE IV, GOR B).</p>
<p>Practice points</p>	<p>Australian NHMRC Guidelines 2010 position statement¹:</p> <ul style="list-style-type: none"> • All newborn infants should receive vitamin K prophylaxis. • Healthy newborn infants should receive vitamin K₁ either: <ul style="list-style-type: none"> ○ By intramuscular injection of 1 mg (0.1 mL) of Konakion® MM Paediatric at birth. This is the preferred route for reliability of administration and level of compliance OR ○ Three 2 mg (0.2 mL) oral doses of Konakion® MM Paediatric, given at: <ul style="list-style-type: none"> ▪ 1st dose: at birth ▪ 2nd dose: 3-5 days of age ▪ 3rd dose: in the fourth week if the baby is fully breastfed. (Babies fed mainly by formula do not need the third dose). ▪ If the infant vomits within one hour of swallowing vitamin K, infant will need to have another dose. • Newborns who are too unwell and are unable to take oral vitamin K₁ (or whose mothers have taken medications that interfere with vitamin K metabolism) should be given 1 mg of Konakion® MM Paediatric by intramuscular injection at birth. A smaller intramuscular dose of 0.5 mg (0.05 mL) should be given to infants with a birth weight of less than 1.5 kg.

	<p>European Society for Paediatric Gastroenterology, Hepatology and Nutrition 2016 position statement⁹</p> <ul style="list-style-type: none"> • Healthy newborn infants should either receive 1 mg of vitamin K₁ by intramuscular injection at birth; or 3 × 2 mg vitamin K₁ orally at birth, at 4 to 6 days and at 4 to 6 weeks; or 2 mg vitamin K₁ orally at birth, and a weekly dose of 1 mg orally for 3 months. • Intramuscular application is the preferred route for efficiency and reliability of administration. • The success of an oral policy depends on compliance with the protocol and this may vary between populations and healthcare settings. <p>American Academy of Pediatrics 2022 Policy statement¹⁰</p> <ul style="list-style-type: none"> • Infants weighing >1500g at birth - A single dose of intramuscular vitamin K 1 mg should be administered within 6 hours of birth. • Infants weighing ≤1500g - A single dose of intramuscular vitamin K 0.3 mg/kg to 0.5 mg/kg should be administered. A single intravenous dose of vitamin K for preterm infants is not recommended for prophylaxis. • VKDB should be considered when evaluating bleeding in the first 6 months of life, even in infants who received prophylaxis, and especially in exclusively breastfed infants.
References	<ol style="list-style-type: none"> 1. 2010 NHMRC Joint statement and recommendations on vitamin K administration to newborn infants to prevent vitamin K deficiency bleeding in infancy (Joint Statement). October 2010. Accessed on 4 April 2021. 2. Puckett RM, Offringa M. Prophylactic vitamin K for vitamin K deficiency bleeding in neonates. Cochrane Database of Systematic Reviews. 2000(4):CD002776. 3. Ardell S, Offringa M, Ovelman C, Soll R. Prophylactic vitamin K for the prevention of vitamin K deficiency bleeding in preterm neonates. Cochrane Database of Systematic Reviews. 2018;2:CD008342. 4. Williams MD, Chalmers EA, Gibson BE. The investigation and management of neonatal haemostasis and thrombosis. British journal of haematology. 2002;119(2):295-309. 5. Raith W, Fauler G, Pichler G, Muntean W. Plasma concentrations after intravenous administration of phylloquinone (vitamin K₁) in preterm and sick neonates. Thrombosis research. 2000;99(5):467-72. 6. Stoeckel K, Joubert P, Grüter J. Elimination half-life of vitamin K 1 in neonates is longer than is generally assumed: implications for the prophylaxis of haemorrhagic disease of the newborn. European journal of clinical pharmacology. 1996;49(5):421-3. 7. Marinova M, Lütjohann D, Breuer O, Kölsch H, Westhofen P, Watzka M, et al. VKORC1-dependent pharmacokinetics of intravenous and oral phylloquinone (vitamin K₁) mixed micelles formulation. European journal of clinical pharmacology. 2013;69(3):467-75. 8. Merative™ Micromedex® Complete IV Compatibility (electronic version). Merative, Ann Arbor, Michigan, USA. Available at: https://www.micromedexsolutions.com/ (cited: Sep/04/2026). 9. ESPGHAN Committee on Nutrition, Mihatsch WA, Braegger C, Bronsky J, Campoy C, Domellöf M, Fewtrell M, Mis NF, Hojsak I, Hulst J, Indrio F. Prevention of vitamin K deficiency bleeding in newborn infants: a position paper by the ESPGHAN committee on nutrition. Journal of pediatric gastroenterology and nutrition. 2016 Jul;63(1):123-9. 10. Hand I, Noble L, Abrams SA; AAP Committee on Fetus and Newborn, Section on Breastfeeding, Committee on Nutrition. Vitamin K and the Newborn Infant. Pediatrics 2022;149(3):e2021056036. 11. Zurynski Y, Grover CJ, Jalaludin B, Elliott EJ. Vitamin K deficiency bleeding in Australian infants 1993–2017: an Australian Paediatric Surveillance Unit study. Archives of Disease in Childhood. 2020 May;105(5):433-8. 12. Simatou E, Tsamantioti E, Hallström A, Stephansson O, Razaz N, Persson M, Bolk J. Vitamin K Prophylaxis in Newborns and Bleeding in Infancy. JAMA pediatrics. 2026.

VERSION/NUMBER	DATE
Original 1.0	3/03/2016
Version 2.0	8/04/2021

Vitamin K₁ (Phytomenadione)

Newborn use only

2026

Version 2.1	12/04/2021
Version 3.0	21/07/2022
Version 3.0 (Minor errata)	10/08/2023
Current 4.0	29/10/2026
REVIEW	29/10/2031

Authors Contribution

Original author/s	Srinivas Bolisetty, Kerrie Knox
Evidence Review	Srinivas Bolisetty, Amber Seigel
Nursing Review	Benjamin Emerson-Parker
Pharmacy Review	Kerrie Knox, Celia Cunha da Silva
ANMF Group contributors	Nilkant Phad, Bhavesh Mehta, Amber Seigel, Jutta van den Boom, Rebecca Barzegar, Mohammad Irfan Azeem, Celia Cunha da Silva, Cindy Chen, Thao Tran, Gloria Yoo, Bryony Malloy, Dianne Lee, Simarjit Kaur, Michelle Jenkins, Renae Gengaroli, Marguerite Mason-Elliott
Final editing	Srinivas Bolisetty
Facilitator	Srinivas Bolisetty

Citation

Australasian Neonatal Medicines Formulary (ANMF). Vitamin K₁ (Phytomenadione). Version number: 4.0. Date of publication 29/10/2026. <https://www.anmfonline.org/>