

<b>Alert</b>	
<b>Indication</b>	Toxoplasmosis
<b>Action</b>	Pyrimethamine is an inhibitor of the enzyme dihydrofolate reductase (DHFR). It blocks the reduction of dihydrofolic acid to tetrahydrofolic acid, leading to disruption of protein synthesis and nuclear division. The affinity of pyrimethamine for protozoal DHFR is much greater than that for the mammalian enzyme. Sulphonamides act synergistically with pyrimethamine by arresting the production of dihydrofolic acid from para-aminobenzoic acid. This results in sequential blockade of the folate pathway of <i>Toxoplasma gondii</i> . (3)
<b>Drug type</b>	Antiprotozoal
<b>Trade name</b>	Daraprim ( <a href="#">Special Access Scheme</a> )
<b>Presentation</b>	Daraprim 25 mg tablets 2 mg/mL oral suspension can be prepared by pharmacy
<b>Dose</b>	<b>Treatment for Congenital Toxoplasmosis</b> Anti-toxoplasma therapy is for <b>12 months</b> and as follows: (1,2) <b>Pyrimethamine</b> First 2 days: 1 mg/kg/dose every 12 hours followed by From Day 3 to 6 months: 1 mg/kg/dose once daily followed by 7 <sup>th</sup> month to 12 months: 1 mg/kg/dose three times a week. <b>Sulfadiazine</b> 50 mg/kg/dose every 12 hours from day 1 of treatment to 12 months and, <b>Calcium folinate (folinic acid)</b> 10 mg three times a week for 12 months until 1 week following cessation of pyrimethamine treatment.
<b>Dose adjustment</b>	Therapeutic hypothermia – Not applicable. ECMO – Not applicable. Renal impairment – Limited data. Caution may be required. (3) Hepatic impairment - Caution is required. (3)
<b>Maximum dose</b>	25 mg/dose
<b>Total cumulative dose</b>	-
<b>Route</b>	Oral
<b>Preparation</b>	<u>2 mg/mL oral suspension prepared in-house by pharmacy:</u> A 2 mg/mL oral suspension may be made with tablets and a 1:1 mixture of Simple Syrup, NF and methylcellulose 1%. Crush forty 25 mg tablets in a mortar and reduce to a fine powder. Add small portions of vehicle and mix to a uniform paste; mix while adding the vehicle in incremental proportions to almost 500 mL; transfer to a calibrated bottle, rinse mortar with vehicle, and add a quantity of vehicle sufficient to make 500 mL. Label "shake well" and "refrigerate". Stable for 91 days.(9,10)
<b>Administration</b>	Give with food/feed Pyrimethamine should be given concurrently with sulfadiazine.(3)
<b>Monitoring</b>	Full blood count especially for neutropenia. Monitoring twice weekly is recommended
<b>Contraindications</b>	History of hypersensitivity to pyrimethamine or any of the components of the preparation.
<b>Precautions</b>	Depression of haematopoiesis: Can occur in about 25% of patients. All patients receiving pyrimethamine should be given a folate/folate supplement to reduce the risk of bone marrow depression. (3) Seizures: Caution should be exercised in patients with a history of seizures. Hepatic impairment: Liver is the main route of metabolism. Caution is required. Galactosemia and hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption – Daraprim contains lactose. (3) G6PD deficiency: Use with caution in patients with possible G6PD deficiency.
<b>Drug interactions</b>	Pyrimethamine may further depress folate metabolism in patients receiving treatment with other folate inhibitors or agents associated with myelosuppression, including co-trimoxazole, trimethoprim, zidovudine or cytostatic agents (e.g. methotrexate). (3) The concurrent administration of lorazepam and pyrimethamine may induce hepatotoxicity. Highly protein bound compounds: Pyrimethamine may displace other drugs from protein binding sites, causing elevated concentrations of unbound drugs; this may be of particular clinical concern for drugs that are both highly protein bound and have a narrow therapeutic index. (3)

<b>Adverse reactions</b>	Hematologic: Leucopenia, anaemia, thrombocytopenia, pancytopenia, neutropenia (4,5) Nervous system: Headache, dizziness, convulsions (6) Gastrointestinal: Anorexia, glossitis (atrophic), vomiting, diarrhoea Dermatologic: Rash (3)
<b>Compatibility</b>	Not applicable.
<b>Incompatibility</b>	Not applicable.
<b>Stability</b>	Oral suspension: 91 days when stored in the fridge.
<b>Storage</b>	Tablets: store below 30°C. Protect from light. Oral suspension: 2-8°C.
<b>Excipients</b>	Lactose, maize starch, hydrolysed starch, docusate sodium and magnesium stearate.(3)
<b>Special comments</b>	-
<b>Evidence</b>	<p><b>Efficacy</b></p> <p><u>Neonates with Congenital Toxoplasmosis:</u> Treatment with the following medications is recommended for 12 months: Pyrimethamine: 1 mg/kg every 12 hours for 2 days followed by 1 mg/kg daily for 6 months followed by the same dose, 3 times a week to complete 12 months; Sulfadiazine: 50 mg/kg every 12 hours; and Folinic acid: 10 mg 3 times a week for 12 months. Folinic acid should be administered until 1 week following cessation of pyrimethamine. (1,2)</p> <p>The United States data suggest that risk of recurrent eye disease is around 31% in infants with CT who had received 12 months of postnatal treatment during their first year of life. (7) The French cohort study showed the risk of recurrence of eye disease within 12 years after the diagnosis of the first eye lesion was around 34%. The French cohort had mothers who were treated during pregnancy and infants who were also postnatally treated. (8)</p> <p><u>Older children (diagnosed beyond neonatal age) with active disease (Chorioretinitis): (1)</u> Treatment is given for at least 1–2 weeks after resolution of all signs and symptoms of acute chorioretinitis (with sharpening of the lesion borders and/or scarring of the lesion) and for ~4–6 weeks total. Acute eye disease often resolves within 10 to 14 days after initiation of treatment, but there are cases that take a longer time to resolve.</p> <p><u>Pyrimethamine</u> First 2 days: 1 mg/kg/dose orally twice a day (maximum 50 mg/day) Then: 1 mg/kg/dose orally once daily (maximum 25 mg/day)</p> <p><u>Sulfadiazine</u> 75 mg/kg/dose orally × 1, followed by 50 mg/kg/dose orally twice a day</p> <p><u>Folinic acid</u> 10–20 mg orally three times a week</p> <p><u>Prednisone (severe chorioretinitis)</u> 0.5 mg/kg/dose twice a day (maximum 40 mg/day; rapid taper)</p> <p><b>Pharmacokinetics</b> It is 87% protein bound. The main route of metabolism is in liver. Plasma half-life is 80-90 hours. Kidney is not the major route of excretion and excretion is not significantly altered in renal failure. (3) Plasma concentration of pyrimethamine were measured in 24 infants aged 1–5 months, treated for congenital toxoplasmosis. Pyrimethamine was used in a single daily dose of 0.35–0.98 mg/kg daily. Concentration of pyrimethamine ranged from 0.01 to 1.2 µg/mL. In 14 children, concentration achieved therapeutic value. In 7 patients, concentration was below therapeutic level, and in 3 patients above therapeutic level. Transient moderate neutropenia was noted in 46% of infants. Modification of therapy was necessary in 12 patients. (11) ANMF consensus: Routine monitoring of plasma concentrations is not practised in Australia.</p> <p><b>Safety</b> Pyrimethamine/sulfadiazine treatment in children was associated with adverse events in 14%-50%.(4,5) Main adverse effect was neutropenia, more often with higher doses and especially when folinic acid was not administered. Seizures have been reported with pyrimethamine overdose. (6)</p>
<b>Practice points</b>	

References	
	1. Maldonado YA, Read JS, Committee On Infectious D. Diagnosis, Treatment, and Prevention of Congenital Toxoplasmosis in the United States. <i>Pediatrics</i> . 2017;139.
	2. Management of Perinatal Infections. Australasian Society for Infectious Diseases (ASID). 2014.
	3. Daraprim. Product information. Accessed on 23 November 2021.
	4. Daveluy A, FH, Bricout H, et al; European Toxoprevention Study Group (EUROTOXO). Review of data related to side effects of drugs used in congenital toxoplasmosis. Panel 2: Treatment Issues. August 24, 2005.
	5. Schmidt DR, Høgh B, Andersen O, Hansen SH, Dalhoff K, Petersen E. Treatment of infants with congenital toxoplasmosis: tolerability and plasma concentrations of sulfadiazine and pyrimethamine. <i>Eur J Pediatr</i> 2006;165(1):19–25.
	6. Genuini M, Freihuber C, Girard I, de Montgolfier I, Kieffer F, Mitancher D. Neonatal intoxication with pyrimethamine: risk due to the absence of pediatric formulation? [in French]. <i>Arch Pediatr</i> 2011;18(10):1084–1086.
	7. Phan L, Kasza K, Jalbrzikowski J, Noble AG, Latkany P, Kuo A, Mieler W, Meyers S, Rabiah P, Boyer K, Swisher C. Longitudinal study of new eye lesions in treated congenital toxoplasmosis. <i>Ophthalmology</i> . 2008 Mar 1;115(3):553-9.
	8. Wallon M, Garweg JG, Abrahamowicz M, et al. Ophthalmic outcomes of congenital toxoplasmosis followed until adolescence. <i>Pediatrics</i> . 2014 Mar 1; 133(3):e601-8.
	9. Nahata MC, Morosco RS, Hipple TF. Stability of pyrimethamine in a liquid dosage formulation stored for three months. <i>American journal of health-system pharmacy</i> . 1997 Dec 1;54(23):2714-6.
	10. Nahata MC, Pai VB, and Hipple TF, <i>Pediatric Drug Formulations</i> , 5th ed, Cincinnati, OH: Harvey Whitney Books Co, 2004.
	11. Lipka B, Milewska-Bobula B, Filipek M. Monitoring of plasma concentration of pyrimethamine (PYR) in infants with congenital <i>Toxoplasma gondii</i> infection—own observations. <i>Wiadomości parazytologiczne</i> . 2011;57(2).

VERSION/NUMBER	DATE
Original	15/02/2022
REVIEW	15/02/2027

**Authors Contribution**

Original author/s	Srinivas Bolisetty, David Osborn
Evidence Review	Srinivas Bolisetty, Tony Lai, Pamela Palasanthiran, Karel Allegaert
Expert review	Tony Lai, Pamela Palasanthiran, Karel Allegaert
Nursing Review	Eszter Jozsa, Sarah Neale, Priya Govindaswamy, Kirsty Minter
Pharmacy Review	Mohammad Irfan Azeem, Joanne Malloy
ANMF Group contributors	Nilkant Phad, Bhavesh Mehta, John Sinn, Cindy Chen, Michelle Jenkins, Helen Huynh, Hannah Bell, Simarjit Kaur, Thao Tran
Final editing and review	Thao Tran
Electronic version	Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty