

Micafungin

Newborn use only

2023

Alert	The Antimicrobial Stewardship Team has listed this drug under the following categories: Restricted (Red category) Use in consultation with a paediatric infectious diseases physician. Not recommended routinely as the first line therapy in neonates with disseminated candidiasis
Indication	Invasive neonatal candidiasis
Action	Selectively inhibits enzymes essential for fungal cell wall synthesis
Drug type	Echinocandin antifungal
Trade name	Mycamine
Presentation	Micafungin Sodium 50mg and 100mg vials
Dose	Invasive candidiasis where central nervous system and/or ocular infection is present or not excluded ^{1,2,3} 10mg/kg/day once a day Invasive candidiasis without meningoencephalitis and/or ocular dissemination ¹ 4mg/kg/day once a day
Dose adjustment	Therapeutic hypothermia: Limited data ECMO: Supplemental dose not required ^{1,4,5} Renal impairment: No dosage adjustment necessary ^{1,6} Hepatic impairment: No dosage adjustment necessary in mild to moderate impairment. ^{1,7}
Maximum dose	15mg/kg/day
Total cumulative dose	
Route	Intravenous
Preparation	Add 5 mL of 5% glucose or 0.9% sodium chloride to 50mg vial to make a concentration of 10 mg/mL. Gently dissolve micafungin powder by swirling the vial to avoid shaking and excessive foaming. Further dilution Peripheral intravenous venous cannula Draw up 5 mL (50mg of micafungin) of the above solution and add to 28 mL of 5% glucose or 0.9% sodium chloride to make a final concentration of 1.5mg/mL. Central venous catheter Draw up 5 mL (50mg of micafungin) of the above solution and add to 7.5 mL of 5% glucose or 0.9% sodium chloride to make a final concentration of 4mg/mL.
Administration	IV infusion: over 1 hour. Cover infusion bag/bottle with an opaque bag to protect from light. More rapid infusions may result in a higher incidence of histamine-mediated reactions. For final concentration greater than 1.5 mg/mL, a central catheter is preferred.
Monitoring	Serum electrolytes, liver and renal function. Monitor IV site for signs of irritation/extravasation.
Contraindications	Hypersensitivity to micafungin, any components of the product or other echinocandins
Precautions	Haemolytic anaemia, thrombocytopenia, moderate to severe hepatopathy, and renal impairment
Drug interactions	Concurrent use may increase plasma levels of itraconazole, nifedipine, methotrexate, and sirolimus.
Adverse reactions	Hyponatremia, hypochloremia, hypokalemia, monocytosis, infusion site phlebitis, fever, rash and transient elevation of hepatic enzymes, diarrhoea, vomiting, elevated creatinine levels, anaemia, and thrombocytopenia. Infusion reactions and anaphylaxis have been reported.
Compatibility	Fluids Sodium chloride 0.9%, 5% glucose, Ringer lactate, and lipid emulsion. Refer to Micromedex for amino acid compatibility. Y-site Aminophylline, calcium chloride, calcium gluconate, dopamine hydrochloride, esmolol, furosemide, heparin, lidocaine, lorazepam, magnesium sulfate, milrinone lactate, glyceryl trinitrate, noradrenaline bitartrate, potassium chloride, sodium nitroprusside, sodium phosphate, tacrolimus, and vasopressin
Incompatibility	Fluids: no information Adrenaline, albumin, amiodarone, dobutamine, insulin, labetalol, midazolam hydrochloride, morphine sulfate, octreotide acetate, phenytoin sodium, rocuronium bromide, and vecuronium bromide
Stability	Infusion solution is stable for 24 hours below 25°C or at 2 to 8°C. Protect from light. ²⁴
Storage	Unopened vial: store below 25°C. ²⁵ Reconstituted solution in vial: can be stored for up to 24hours below 25°C, protected from light. ²⁵

	Protect from light.
Excipients	Lactose, citric acid, and sodium hydroxide.
Special comments	May cause fever, and injection-site pain and inflammation. Anaphylactic reactions have been reported.
Evidence	<p>Efficacy</p> <p>Invasive candidiasis is reported in 1.5% to 9% very low birth weight neonates.^{8,9} It is associated with high mortality (20-40%) and neurodevelopmental disability (50-60%) in survivors.¹⁰</p> <p>Treatment of candida infection</p> <p>In one RCT of 98 children which included 7 neonates, micafungin was administered for a median duration of 14 days to treat acute candidaemia. The treatment was commenced at a daily dose of 2mg/kg/day and increased to 4mg/kg/day if clinical signs of candidaemia or positive blood culture persisted for 5 days or more. In this trial the rate of overall treatment success for micafungin (73%) was comparable with liposomal amphotericin-B (76%). During the entire study including the 12-week follow-up, the mortality rates were 25% and 24% for the micafungin and amphotericin-B group respectively. 4% of participants from the micafungin group and 16% from the liposomal amphotericin-B group had treatment discontinued due to adverse events.¹¹ In another RCT, 20 infants aged < 4 months received 10mg/kg/day micafungin for a mean duration of 18 days for proven invasive candida infection.¹² <i>Candida albicans</i> (41%) and <i>Candida parapsilosis</i> (34%) were the most commonly isolated organisms. A complete clinical response and fungal free survival at 1 week after completion of the treatment was achieved in 60% participants. 25% of participants were alive after 1 week of treatment but were not free of fungal infection. Authors reported recurrence of infection in about 10-15% patient receiving micafungin.^{11,12} Auriti et al used micafungin at a dose of 8-15mg/kg/day in a cohort of 18 infants with invasive candida infection including meningitis. The mean gestational age of the infants at birth was 35 weeks and the mean postnatal age was 9 weeks at treatment. <i>Candida</i> was isolated from blood in 10, urine in 6, and CSF in 2 participants. The mean duration of treatment for <i>C. Albicans</i> (n=9) and <i>C. Parapsilosis</i> (n=7) was 19 days. The overall survival with clinical resolution of infection was 78%. In this study 20% of the surviving infants developed neurological impairment.¹³</p> <p>Prophylaxis for fungal infection</p> <p>When co-administered with enteral miconazole, micafungin at a dose of 3mg/kg/day intravenously was more effective than fluconazole in prevention of fungal infection (71 vs 39% success) in extremely preterm infants.^{14,15} In a cohort study primarily designed to study the pharmacokinetics of prophylactic dose, 25 very low birthweight who received intravenous micafungin at a dose of 1mg/kg/day for a mean duration of 13 days did not develop fungal infection.¹⁸</p> <p>ANMF consensus: Prophylaxis with micafungin should only be given in consultation with a paediatric ID specialist, in situations where prophylaxis is indicated and neither nystatin nor fluconazole are feasible/appropriate.</p> <p>Safety</p> <p>Although results from several trials of micafungin in infants have shown the drug to be generally well tolerated, these trials are small. Studies have reported reversible haemolytic anaemia in as high as 30-40% of the recipients. Moderate elevation of serum creatinine, 2-3 times higher levels of hepatic transaminases and disturbances in serum electrolytes have also been reported in up to 20-40% participants. Thrombocytopenia has been described in about 10% patients receiving micafungin.^{12,16}</p> <p>Pharmacokinetics</p> <p>Several studies have assessed the pharmacokinetics of micafungin in neonates.^{1,12,17-22} Micafungin displays linear pharmacokinetics with minimal systemic accumulation after repeated administration. A steady state is reached within 4 days with once daily dosing regimen. Its volume of distribution (0.39–0.76L/kg) in premature neonates is comparable to adults. In neonates, 96.7% micafungin is bound to albumin and its elimination half-life is (5.5–11 hours) shorter than adults. Total body clearance of micafungin is faster in neonates compared to children and adults. Micafungin concentrates the highest in the lungs, liver, spleen, and kidneys. It is metabolized primarily by the liver into various metabolites and excreted in bile. Faecal excretion is the major route of elimination and < 1% of micafungin is excreted unchanged in the urine. Micafungin has poor oral bioavailability.</p> <p>Hope et al pooled the pharmacokinetic data on micafungin for suspected or proven invasive candida infection on 47 infants aged < 4 months across three trials.^{2,12,17,19} In these trials micafungin was administered at a dose ranging from 0.75 to 15mg/kg/day. The proportion of the patients receiving 8, 10, and 12mg/kg/day with an AUC less than the target (166.5mg x h/L) was 29.3%, 17.4%, and 10.5%, respectively. A dosage of 10mg/kg/day resulted in 82.6% of patients with AUCs that are associated with near-maximal decline in fungal burden within the central nervous system. The AUCs also suggested that</p>

	at 15mg/kg/day micafungin in preterm neonates provided a similar systemic exposure to a dose of approximately 5mg/kg in adults. Based on Monte Carlo simulations of 81 concentrations, Leroux et al showed attainment of the AUC ₀₋₂₄ target of 166.5mg x h/L in 100% neonates when micafungin was administered at a dose of 10mg/kg/day with or without a 15mg/kg loading dose. ²³
Practice points	
References	<ol style="list-style-type: none"> 1. Abdel-Haq N, Smith SM, Asmar BI. Micafungin injection for the treatment of invasive candidiasis in pediatric patients under 4 months of age. <i>Expert Rev Anti Infect Ther.</i> 2022 Apr; 20(4):493-505 2. Hope WW, Smith PB, Arrieta A, et al. Population pharmacokinetics of micafungin in neonates and young infants. <i>Antimicrob Agents Chemother.</i> 2010 Jun; 54(6):2633-7 3. Ascher S, Smith PB, Benjamin DK Jr. Safety of micafungin in infants: insights into optimal dosing. <i>Expert Opin Drug Saf.</i> 2011 Mar; 10(2):281-6 4. Autmizguine J, Hornik CP, Benjamin DK Jr, et al. Pharmacokinetics and Safety of Micafungin in Infants Supported With Extracorporeal Membrane Oxygenation. <i>Pediatr Infect Dis J.</i> 2016 Nov; 35(11):1204-1210. 5. Sherwin J, Heath T, Watt K. Pharmacokinetics and Dosing of Anti-infective Drugs in Patients on Extracorporeal Membrane Oxygenation: A Review of the Current Literature. <i>Clin Ther.</i> 2016 Sep; 38(9):1976-94. 6. Greenberg RG, Benjamin DK Jr. Neonatal candidiasis: diagnosis, prevention, and treatment. <i>J Infect.</i> 2014 Nov; 69 Suppl 1(0 1):S19-22 7. Wasmann RE, Muilwijk EW, Burger DM, et al. Clinical Pharmacokinetics and Pharmacodynamics of Micafungin. <i>Clin Pharmacokinet.</i> 2018 Mar; 57(3):267-286 8. Benjamin DK Jr, Stoll BJ, Gantz MG, et al. Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Neonatal candidiasis: epidemiology, risk factors, and clinical judgment. <i>Pediatrics.</i> 2010 Oct; 126(4):e865-73. 9. Barton M, O'Brien K, Robinson JL, et al. Invasive candidiasis in low birth weight preterm infants: risk factors, clinical course and outcome in a prospective multicenter study of cases and their matched controls. <i>BMC Infect Dis.</i> 2014 Jun 12; 14:327. 10. Friedman S, Richardson SE, Jacobs SE, O'Brien K. Systemic Candida infection in extremely low birth weight infants: short term morbidity and long term neurodevelopmental outcome. <i>Pediatr Infect Dis J.</i> 2000 Jun; 19(6):499-504 11. Queiroz-Telles F, Berezin E, Leverger G, et al. Micafungin Invasive Candidiasis Study Group. Micafungin versus liposomal amphotericin B for pediatric patients with invasive candidiasis: substudy of a randomized double-blind trial. <i>Pediatr Infect Dis J.</i> 2008 Sep; 27(9):820-6 12. Benjamin DK Jr, Smith PB, Arrieta A, et al. Safety and pharmacokinetics of repeat-dose micafungin in young infants. <i>Clin Pharmacol Ther.</i> 2010 Jan; 87(1):93-9. 13. Auriti C, Goffredo BM, Ronchetti MP, et al. High-Dose Micafungin in Neonates and Young Infants with Invasive Candidiasis: Results of a Phase 2 Study. <i>Antimicrob Agents Chemother.</i> 2021 Mar 18; 65(4):e02494-20. 14. Maede Y, Ibara S, Nagasaki H, et al. Micafungin versus fluconazole for prophylaxis against fungal infections in premature infants. <i>Pediatr Int.</i> 2013 Dec; 55(6):727-30. 15. Ferrando G, Castagnola E. Prophylaxis of Invasive Fungal Infection in Neonates: A Narrative Review for Practical Purposes. <i>J Fungi (Basel).</i> 2023 Jan 26; 9(2):164 16. Schüller SS, Bauer C, Unterasinger L, Berger A. Safety and Efficacy of Micafungin in Extremely Low Birth Weight Infants. <i>Pediatr Infect Dis J.</i> 2018 Jun; 37(6):e169-e172. 17. Heresi GP, Gerstmann DR, Reed MD, et al. The pharmacokinetics and safety of micafungin, a novel echinocandin, in premature infants. <i>Pediatr Infect Dis J.</i> 2006 Dec; 25(12):1110-5 18. Kawada M, Fukuoka N, Kondo M, et al. Pharmacokinetics of prophylactic micafungin in very low birth weight infants. <i>Pediatr Infect Dis J.</i> 2009 Sep; 28(9):840-2. 19. Smith PB, Walsh TJ, Hope W, et al. Pharmacokinetics of an elevated dosage of micafungin in premature neonates. <i>Pediatr Infect Dis J.</i> 2009 May; 28(5):412-5 20. Manzoni P, Wu C, Tweddle L, et al. Micafungin in premature and non-premature infants: a systematic review of 9 clinical trials. <i>Pediatr Infect Dis J.</i> 2014 Nov; 33(11):e291-8. 21. Yanni SB, Smith PB, Benjamin DK Jr, et al. Higher clearance of micafungin in neonates compared with adults: role of age-dependent micafungin serum binding. <i>Biopharm Drug Dispos.</i> 2011 May; 32(4):222-32 22. De Rose DU, Bersani I, Ronchetti MP, et al. Plasma and Cerebrospinal Fluid Concentrations of Micafungin Administered at High Doses in Critically Ill Infants with Systemic Candidiasis: A Pooled Analysis of Two Studies. <i>Pharmaceuticals (Basel).</i> 2023 Mar 22; 16(3):472

	<p>23. Leroux S, Jacqz-Aigrain E, Elie V, et al. FP7 TINN (Treat Infections in NeoNates) consortium. Pharmacokinetics and safety of fluconazole and micafungin in neonates with systemic candidiasis: a randomized, open-label clinical trial. Br J Clin Pharmacol. 2018 Sep; 84(9):1989-1999</p> <p>24. Briot T, Vrignaud S, Lagarce F. Stability of micafungin sodium solutions at different concentrations in glass bottles and syringes. Int J Pharm Sci. 2015 Aug; 492(1-2):137-40.</p> <p>25. Product information. Available from www.tga.gov.au</p> <p>26. MerativeTM Micromedex® Complete IV Compatibility (electronic version). Merative, Ann Arbor, Michigan, USA. Available at: https://www.micromedexsolutions.com/ (cited: July/24/2023).</p>
--	---

VERSION/NUMBER	DATE
Original	26/07/2023
REVIEW	26/07/2028

Authors Contribution

Original author/s	Nilkant Phad
Evidence Review	Nilkant Phad
Expert review	Tony Lai, Brendan McMullan
Nursing Review	Eszter Jozsa, Ben Emerson-Parker
Pharmacy Review	Stephanie Halena, Michelle Jenkins
ANMF Group contributors	Bhavesht Mehta, Srinivas Bolisetty, Mohammad Irfan Azeem, Rebecca O-Grady, Rebecca Barzegar, Thao Tran, Helen Huynh, Simarjit Kaur, Susannah Brew, Renae Gengaroli, Martin Kluckow, Karel Allegaert
Final editing	Ben Emerson-Parker
Electronic version	Cindy Chen, Ian Callander
Facilitator	Nilkant Phad, Srinivas Bolisetty

Citation for the current version

Phad N, Lai T, McMullan B, Bolisetty S, Jozsa E, Emerson-Parker B, Halena S, Jenkins M, Mehta B, Azeem MI, O'Grady R, Kaur S, Barzegar R, Kluckow M, Tran T, Huynh H, Brew S, Gengaroli R, Chen C, Callander I, Allegaert K. Micafungin. Consensus formulary by the Australasian Neonatal Medicines Formulary group. Version 2, dated 26 July 2023. www.anmfonline.org