

Alert	<p>High risk medication. Also known as unfractionated heparin (UFH). Not equivalent to low molecular weight heparin (LMWH). Use in consultation with haematologist for treatment of thrombosis. Many concentrations of heparin are available. Accidental overdose can occur when multiple concentrations are kept in the unit. In neonatal settings, recommend storing the following preparations only: heparinised saline 50 units/5 mL and heparin sodium injection ampoule 1000 units/1 mL. DBL Heparin sodium injection in vials is not recommended in neonates as it contains benzyl alcohol. However, DBL Heparin sodium injection in ampoules does not contain benzyl alcohol.</p>												
Indication	<p>Primary or secondary antithrombotic prophylaxis. Maintenance of arterial and central venous catheter patency.</p>												
Action	<p>Heparin binds to antithrombin III (ATIII), potentiating ATIII's activity by at least 1000-fold. ATIII predominantly inactivates factor Xa and thrombin (other proteases/clotting factors to lesser degree), which in turn inhibits conversion of fibrinogen to fibrin. Heparin also possesses anti-complementary activity, inhibiting both the classic and alternative pathways.</p>												
Drug type	<p>Anticoagulant</p>												
Trade name	<p>Heparin Sodium Injection (Pfizer) DBL Heparin Sodium Injection BP Heparinised Saline Injection (Pfizer)</p>												
Presentation	<p>Antithrombotic prophylaxis Pfizer Heparin Sodium Injection Ampoule: 5000 units/5 mL DBL Heparin Sodium Injection BP Ampoule: 1000 units/1 mL DBL Heparin Sodium BP Vials – Not to be used in neonates as it contains benzyl alcohol.</p> <p>Maintenance of catheter patency Heparinised Saline Injection Ampoule: 50 units/5 mL (10 units/mL) Also available as commercial premade 50mL syringe containing 1 unit/mL of heparin in sodium chloride 0.9%.</p>												
Dose	<p>Antithrombotic prophylaxis^{1,2,3} Loading dose: 75 (50-100) units/kg over 30 minutes. Initial maintenance dose: 30 (20-40) units/kg/hour as continuous IV infusion.</p> <p>Adjustment of Heparin dose Anti-Xa is preferred to assess the effect of heparin and guide dosing (Table 1). If anti-Xa levels are not available, APTT (Activated partial thromboplastin time) can be used to guide heparin dosing (Table 2).</p> <p>Table 1. Heparin dosing based on anti-Xa levels (therapeutic range 0.3-0.7 unit/mL) (modified from O'Meara et al)³</p> <table border="1" data-bbox="343 1552 1481 1753"> <thead> <tr> <th>Anti-Xa level (unit/mL)</th> <th>Dose adjustment</th> </tr> </thead> <tbody> <tr> <td><0.2</td> <td>Increase infusion by 5 units/kg/hour</td> </tr> <tr> <td>0.2-0.29</td> <td>Increase infusion by 5 units/kg/hour</td> </tr> <tr> <td>0.3-0.7</td> <td>No change</td> </tr> <tr> <td>0.71-1.0</td> <td>Decrease infusion by 2 unit/kg/hour</td> </tr> <tr> <td>>1.0</td> <td>Seek advice from haematologist</td> </tr> </tbody> </table> <p>Measure anti-Xa levels 6 hours after commencing heparin and then 6 hourly until two consequent values are within therapeutic range. After every heparin adjustment or a blood product administration, the anti-Xa level should be checked again in 6 hours and discuss with haematologist on frequency of further monitoring. PT/INR, PTT, fibrinogen, platelet count, and ATIII levels are measured daily or as advised by the haematologist.</p>	Anti-Xa level (unit/mL)	Dose adjustment	<0.2	Increase infusion by 5 units/kg/hour	0.2-0.29	Increase infusion by 5 units/kg/hour	0.3-0.7	No change	0.71-1.0	Decrease infusion by 2 unit/kg/hour	>1.0	Seek advice from haematologist
Anti-Xa level (unit/mL)	Dose adjustment												
<0.2	Increase infusion by 5 units/kg/hour												
0.2-0.29	Increase infusion by 5 units/kg/hour												
0.3-0.7	No change												
0.71-1.0	Decrease infusion by 2 unit/kg/hour												
>1.0	Seek advice from haematologist												

Table 2. Heparin dosing based on APTT levels (therapeutic range 60-85 seconds).^{1,4}

APTT (seconds)	Bolus (units/kg)	Hold (min)	Rate change (%)	Time until repeat APTT
<50	50	0	+10	6 h
50-59	0	0	+10	6 h
60-85	0	0	No change	Next day or as per haematologist advice
86-95	0	0	-10	6 h
96-120	0	30	-10	6 h
>120	0	60	-10	6 h

Obtain blood for APTT 6 hours after administration of loading dose and 6 hours after every change. When APTT values are therapeutic, blood count and APTT daily or as per the advice of haematologist.

Venous catheter patency maintenance.^{1,2,5-7,17-20}

0.5 unit/mL of heparinised saline to run at 0.5 –1 mL/hour. Refer to evidence section.

Arterial catheter patency maintenance.^{1,2,5-7,17-20}

1 unit/mL of heparinised saline to run at 0.5 – 1 mL/hour. Refer to evidence section.

Heparin Lock for Central Venous Access Device (CVAD)

Heparinised saline **10 units/mL** with **0.5 to 1 mL** instilled per lumen as per the priming volume.

Dose adjustment

Therapeutic hypothermia – No information.
ECMO – Refer to local ECMO protocols for anticoagulation.
Renal impairment – Dose adjustment may be required in severe renal impairment. Discuss with haematologist.
Hepatic impairment – No dose adjustment is required.⁸

Route

IV, intra-arterial

Preparation

Antithrombotic prophylaxis - IV infusion:
Use heparin 1000 units/mL for preparation

20mL Syringe

Draw up heparin and add compatible fluid* as per table below to make a final volume of 20 mL.

Heparin concentration	100 units/mL
Volume of heparin (1000 unit/mL)	2 mL (=2000 units)
Volume of compatible fluid*	18 mL
Total volume	20 mL

* Compatible fluid: Sodium chloride 0.9%

50mL Syringe

Draw up heparin and add compatible fluid* as per table below to make a final volume of 50 mL.

Heparin concentration	100 units/mL
Volume of heparin (1000 unit/mL)	5 mL (=5000 units)
Volume of compatible fluid*	45 mL
Total volume	50 mL

* Compatible fluid: Sodium chloride 0.9%

Catheter patency – Arterial/venous infusion

Use heparinised saline 50 units/5 mL for preparation

Venous catheter patency

Add 25 units (2.5 mL) of **heparinised saline** to 47.5 mL of sodium chloride 0.9% or 0.45% to make a final volume of 50 mL with a concentration of 0.5 unit/mL.

Arterial catheter patency

Add 50 units (5 mL) of **heparinised saline** to 45 mL of sodium chloride 0.9% or 0.45% to make a final volume of 50 mL with a concentration of 1 unit/mL.

Commercial premade 50mL syringe containing 1 unit/mL of heparin in sodium chloride 0.9%.

Administration	<p>Systemic antithrombotic therapy Loading dose: Administer over 30 minutes. Maintenance: Continuous IV infusion.</p> <table border="1" style="width: 100%; border-collapse: collapse; margin: 10px 0;"> <tr> <td style="width: 60%;">Heparin concentration</td> <td style="text-align: center;">100 units/mL</td> </tr> <tr> <td>75 units/kg load over 30 minutes is equal to</td> <td style="text-align: center;">0.75 mL/kg over 30 minutes</td> </tr> <tr> <td>30 units/kg/hour is equal to</td> <td style="text-align: center;">0.3 mL/kg/hour</td> </tr> </table> <p>Vascular catheter patency Continuous IV infusion.</p>	Heparin concentration	100 units/mL	75 units/kg load over 30 minutes is equal to	0.75 mL/kg over 30 minutes	30 units/kg/hour is equal to	0.3 mL/kg/hour				
Heparin concentration	100 units/mL										
75 units/kg load over 30 minutes is equal to	0.75 mL/kg over 30 minutes										
30 units/kg/hour is equal to	0.3 mL/kg/hour										
Monitoring	<p>Antithrombotic prophylaxis Six hours after initiating therapy, measure anti-Xa (or APTT if anti-Xa is not available), then adjust dose to achieve anti-Xa level of 0.3 to 0.7 unit/mL (equivalent to APTT of 60 to 85 seconds) – Refer to tables 1 and 2 in the dosing section. Platelet count before the commencement and then weekly. Assess for signs of bleeding and thrombosis.</p> <p>Vascular catheter patency Standard observations for intravascular catheters.</p>										
Contraindications	<p>Known hypersensitivity to heparin, uncontrolled bleeding. Intraventricular haemorrhage, gastrointestinal haemorrhage, thrombocytopenia $< 50 \times 10^9/L$, severe hypertension. Eye, brain, or spinal cord surgery – Surgeons to give clearance regarding when to start heparin.⁷</p>										
Precautions	<p>Bleeding disorders – Discuss with haematologist. Store heparinised saline ampoules separately from other heparin products and sodium chloride 0.9% ampoules to reduce the risk of selection errors.</p>										
Drug interactions	<p>Paracetamol, non-steroid anti-inflammatory drugs, alprostadil, thrombolytic agents, and vitamin A may increase the risk of bleeding.</p>										
Adverse reactions	<p>Haemorrhage and haematoma formation. Heparin-induced thrombocytopenia (HIT). Osteoporosis. Cholestatic liver reaction and elevation of transaminases. Hyperaldosteronism can occur after prolonged administration.⁸</p> <p>Treatment of Heparin-Induced bleeding: (1) cease heparin and (2) if immediate reversal is required, administer protamine sulfate. The required dose of protamine sulfate is based on the amount of UFH received in the previous 2 hours as follows:¹</p> <table border="1" style="width: 100%; border-collapse: collapse; margin: 10px 0;"> <thead> <tr> <th style="text-align: center;">Time Since Last Heparin Dose</th> <th style="text-align: center;">Protamine dose per 100 units of heparin received in the last 2 hours</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;"><30 min</td> <td style="text-align: center;">1 mg</td> </tr> <tr> <td style="text-align: center;">30-60 min</td> <td style="text-align: center;">0.5-0.75 mg</td> </tr> <tr> <td style="text-align: center;">60-120 min</td> <td style="text-align: center;">0.375-0.5 mg</td> </tr> <tr> <td style="text-align: center;">>120 min</td> <td style="text-align: center;">0.25-0.375 mg</td> </tr> </tbody> </table> <p>Maximum dose of 50 mg. Infusion rate of a 10 mg/mL solution should not exceed 5 mg/min. Hypersensitivity reactions to protamine sulfate may occur in patients with known hypersensitivity reactions to fish or those previously exposed to protamine therapy or protamine-containing insulin. For more information, refer to Protamine formulary.</p>	Time Since Last Heparin Dose	Protamine dose per 100 units of heparin received in the last 2 hours	<30 min	1 mg	30-60 min	0.5-0.75 mg	60-120 min	0.375-0.5 mg	>120 min	0.25-0.375 mg
Time Since Last Heparin Dose	Protamine dose per 100 units of heparin received in the last 2 hours										
<30 min	1 mg										
30-60 min	0.5-0.75 mg										
60-120 min	0.375-0.5 mg										
>120 min	0.25-0.375 mg										
Overdose	<p>Protamine sulfate is the reversal agent to correct the anticoagulant effect of heparin. AUSTRALIA: Contact the Poisons Information Centre on 13 11 26 for information on the management of overdose. NEW ZEALAND: Contact the National Poisons Centre on 0800 764 766 for information on the management of overdose.</p>										
Compatibility	<p>Fluids: Sodium chloride 0.9%⁸ PN at Y-site: Amino acid-glucose solution Y-site: Aciclovir, ampicillin, atropine, aztreonam, caffeine citrate, cefazolin, cefTAZIDIME, cefTRIAZONE sodium, cefotaxime, clindamycin, clonidine hydrochloride, dexAMETHASOne, dexMEDETOMIDine, digoxin, dopamine, ephedrine sulfate, ertapenem sodium, fentanyl, fluconazole, folic acid (sodium salt),</p>										

	Ganciclovir sodium, hydrocortisone sodium succinate, HYDRomorphone hydrochloride, levetiracetam, lidocaine hydrochloride, linCOMYCIN hydrochloride, linEZOLID, magnesium sulfate, meropenem, metronidazole, micafungin sodium, morphine sulfate, naloxone hydrochloride, nitroprusside sodium, noradrenaline, octreotide acetate, pancuronium bromide, paracetamol, penicillin G potassium, penicillin G sodium, phenobarbital sodium, piperacillin/tazobactam, potassium chloride, propofol, rocuronium bromide, sodium bicarbonate, suxamethonium, vecuronium, and zidovudine.
Incompatibility	<p>Fluids: Glucose 5%, glucose 10% - Conflicting published reports on the stability of heparin activity in dextrose-containing infusion solutions exist. However, most research indicates that heparin sodium may be given in dextrose-containing solutions up to a concentration of 40 units/mL.⁸</p> <p>PN at Y-site: Lipid emulsion</p> <p>Y-site: Amiodarone hydrochloride, benzylpenicillin, ciprofloxacin, cisatracurium, DIAzepam, diazoxide, ketamine, levetiracetam, phenytoin sodium, and tobramycin.</p> <p>Variable at Y-site: Amikacin sulfate, amphotericin B deoxycholate, ampicillin sodium, calcium chloride, calcium gluconate, cisatracurium besylate, dobutamine hydrochloride, furosemide, gentamicin sulfate, insulin regular, labetalol hydrochloride, methylprednisolone sodium succinate, midazolam hydrochloride, morphine hydrochloride, pantoprazole sodium, sildenafil citrate, sulfamethoxazole-trimethoprim, tobramycin sulfate, and vancomycin hydrochloride.</p>
Stability	
Storage	Ampoule and vial: Store below 25°C. Bag: Store below 30°C.
Excipients	Pfizer ampoule: Water for injection. DBL ampoule: Hydrochloric acid, sodium hydroxide. DBL vial: Benzyl alcohol. Do not give products that contain benzyl alcohol to neonates. Heparinised saline: Hydrochloric acid, sodium chloride, sodium hydroxide.
Special comments	
Evidence	<p>Efficacy</p> <p><u>Systemic antithrombotic therapy/prophylaxis</u></p> <p><u>Arterial thrombosis:</u> Spontaneous arterial thrombosis is rare in neonates and the evidence around its management using heparin is limited to case reports only. De Godoy et al reported complete disappearance of an aortic thrombus and clinical improvement in a neonate following 15 days anticoagulation with heparin.¹⁰ Similarly, anticoagulation with heparin following initial thrombolysis of a major aortic thrombus is found to be helpful in improving clinical outcomes of neonates.¹¹</p> <p><u>Venous thrombosis:</u> In a cohort of 53 neonates who received heparin, Moharir et al found significant reduction in propagation of cerebral sino-venous thrombosis (2 vs 30%; P < 0.001). However, no difference was noted in thrombus recanalisation, mortality and long-term disability.¹² Non-life threatening bleeding was seen in 5-6% of neonates.</p> <p>In two retrospective reviews involving 100 neonates who received heparin therapy for renal vein thrombosis with or without inferior vena cava involvement, there was no difference in irreversible renal damage and renal atrophy at long term follow up.^{13, 14} In a cohort of 128 neonates with portal vein thrombosis the incidence of lobar atrophy of liver and risk of portal hypertension was not altered by the use of anticoagulants.¹⁵</p> <p>No clinical outcome studies have determined the therapeutic range for heparin in neonates and the APTT therapeutic range and monitoring is extrapolated from adults. One prospective cohort study used a weight-based nomogram to address dosing of heparin in paediatric patients required to achieve adult therapeutic APTT values. Bolus doses of 75 to 100 units/kg resulted in therapeutic APTT values in 90% of children at 4-6 hours after bolus.¹⁶</p> <p><u>Maintenance of patency of central vascular catheters</u>^{1,2, 5-7}</p> <p>Low dose heparin administered as a continuous infusion or regular flushes significantly increases the duration of peripheral catheter patency and reduces the episodes of infusion failure.^{5,6} A systematic review involving 267 neonates reported significant reduction in occlusion of peripherally placed percutaneous central venous catheters and higher rates of completion of therapy if heparin is infused at a dose of 0.5 unit/kg/h.⁷ Administration of heparin in low doses does not significantly alter the risk of sepsis or intraventricular haemorrhage.^{1,5-7} However, Lesko et. al. reported a 4-fold, but statistically not significant, increase in IVH in low-birthweight infants in a case control study (OR, 3.9; 95% CI, 1.4-11.0).⁹</p> <p><u>Maintenance of patency of peripheral arterial catheters</u></p>

	<p>Heparin is shown to significantly reduce clot formation and maintain patency of peripheral arterial catheter for a longer period.¹⁷ Compared with 1 unit/mL, heparin concentration of 5 units/mL is more effective in keeping arterial catheters patent for longer duration.¹⁸ Studies found heparinised normal saline superior to heparinised glucose solution, and continuous infusion of heparin in normal saline better compared to intermittent flushing to improve arterial catheter patency.^{19,20}</p> <p>ANMF consensus on the strength and infusion rate of heparinised IV fluid for catheter patency is a simplified pragmatic recommendation from the evidence.</p> <p><u>Heparin Lock for Central Venous Access Device (CVAD)</u></p> <p>The 'lock' is the intraluminal injection of a limited volume of fluid, after the catheter flush, in the intervals of time when the catheter is not in use, with the purpose of preventing lumen occlusion and/or bacterial colonization. The most appropriate lock solution for central venous access devices is still to be defined. The data available from the literature are still not conclusive and no recommendation is offered by most guidelines.²⁴ The recommended regimen suggested by the ANMF group is reflective of the current practice in neonatal units.</p> <p>Safety</p> <p>Major bleeding has been reported in children treated for deep vein thrombosis/pulmonary embolism. There are case reports of osteoporosis. Given the adverse effects, and the availability of alternative anticoagulants, long term use of heparin can be avoided. Heparin-induced thrombocytopenia (HIT) has been reported in neonates. Following exposure to heparin for at least 5 days, Schmutz et al reported antibodies against HPF4 in 2.3% children who developed thrombocytopenia and thrombosis.²² In a systematic review, Avila et. al. reported seroconversion for anti-PF4/H antibodies in 0-1.7% neonates but no neonate fulfilled the combined clinical and laboratory criteria used for the diagnosis of HIT.²³</p> <p>Pharmacokinetics</p> <p>Studies of heparin in newborns are limited but show that the clearance is faster than for older children because of a larger volume of distribution. It is metabolised by liver and excreted renally within 6 hours but may be delayed. Half-life is dose-dependent but averages 1 to 3 hours. Efficacy in neonates may be low due to low antithrombin plasma concentrations.¹</p>
<p>Practice points</p>	<p>General</p> <p>There are no data from randomised controlled trials to recommend or refute the use of heparin for treatment of neonatal thrombosis.²</p> <p>Dose</p> <p><u>Antithrombotic prophylaxis</u></p> <p>Loading doses and maintenance doses have been adapted from the American College of Chest Physicians Evidence-Based Clinical Practice Guidelines 2012,¹ which were based on paediatric data from a prospective cohort study.²¹ (LOE IV GOR D)</p> <p>Loading dose is safer to be infused over 30 minutes in neonates. (ANMF haematology expert group opinion)</p> <p>Initial maintenance dose is easier to be administered at 30 units/kg/h, rather than 28 units/kg/h. (ANMF haematology expert group opinion)</p> <p><u>Central vascular catheters</u></p> <p>Heparin infusions at 0.5 units/kg per hour are recommended to maintain CVAD patency.^{1,7} (LOE I, GOR B)</p> <p><u>Peripheral arterial catheters</u></p> <p>Heparin infusions at 0.5 units/mL at 1 mL/hour are recommended.¹ (LOE II, GOR B)</p> <p>ANMF consensus on the strength and infusion rate of heparinised IV fluid for catheter patency is a simplified pragmatic recommendation from the evidence.</p> <p><u>Heparin Lock for Central Venous Access Device (CVAD)</u></p> <p>The recommended regimen suggested by the ANMF group is reflective of the current practice in neonatal units.</p> <p>Dose adjustment</p> <p>Anti-Xa therapeutic range: While O'Meara study suggests 0.4 – 0.8 unit/mL, range of 0.3 – 0.7 unit/mL is adequate for most indications, and most commonly used. Table 1 is a modified regimen of O'Meara study,³ which was performed in ECMO patients where very tight anticoagulation is required, managed by staff very experience in managing anticoagulation for ECMO circuits; hence, the repeat boluses were recommended by O'Meara et. al. when anti-Xa was below the target range. Repeat boluses are not required in the majority of non-ECMO patients. Regarding dose adjustment for anti-Xa > 1, advice from the</p>

	<p>haematologist should be sought as the anti-Xa can be very high and simply reducing the infusion rate may not be appropriate.³ (ANMF haematology expert group opinion)</p> <p>The frequency of testing at 2 hourly intervals is the practice in ECMO circuits but not indicated for routine anti-coagulation for non-ECMO patients. Testing too early & too frequently, lends to inappropriate dose adjustments. Testing 6 hours after starting infusion and dose changes is adequate as a general guide, and to check with the haematologist on further monitoring. (ANMF haematology expert group opinion)</p> <p>Dose adjustments using APTT monitoring have been adapted from the American College of Chest Physicians Evidence-Based Clinical Practice Guidelines 2012,¹ which were based on paediatric data from a prospective cohort study.²¹ (LOE IV GOR D)</p> <p>For consistency, using APTT monitoring, testing 6 hours after starting infusion and dose changes is suggested as a general guide, and to check with the haematologist. (ANMF haematology expert group opinion)</p>
<p>References</p>	<ol style="list-style-type: none"> 1. Monagle P, Chan A, Goldenberg N et al: Antithrombotic Therapy in Neonates and Children: Antithrombotic Therapy and Prevention of Thrombosis, 9th Ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. <i>Chest</i> Feb, 2012; 141(2 suppl): e737S-e801S. 2. Romantsik O, Bruschetti M, Zappettini S, et al. Heparin for the treatment of thrombosis in neonates. <i>Cochrane Database Syst Rev</i>. 2016 Nov 7; 11:CD012185. 3. O'Meara LC, Alten JA, Goldberg KG, Timpa JG, Phillips J, Laney D, Borasino S. Anti-xa directed protocol for anticoagulation management in children supported with extracorporeal membrane oxygenation. <i>ASAIO J</i>. 2015 May-Jun;61(3):339-44 4. Bhatt MD, Paes BA, Chan AK. How to use unfractionated heparin to treat neonatal thrombosis in clinical practice. <i>Blood Coagul Fibrinolysis</i>. 2016 Sep;27(6):605 5. Upadhyay A, Verma K, Lal P, et al. Heparin for prolonging peripheral intravenous catheter use in neonates: a randomized controlled trial. <i>J Perinatol</i>. 2015 Apr; 35(4):274-7. 6. Kumar M, Vandermeer B, Bassler D, Mansoor N. Low-dose heparin use and the patency of peripheral IV catheters in children: a systematic review. <i>Pediatrics</i>. 2013 Mar;131(3):e864-72 7. Shah P, Shah V. Continuous heparin infusion to prevent thrombosis and catheter occlusion in neonates with peripherally placed percutaneous central venous catheters. <i>Cochrane Database Syst Rev</i>. 2008 Apr 16 ;(2):CD002772. 8. Merative™ Micromedex® Complete IV Compatibility (electronic version). Merative, Ann Arbor, Michigan, USA. Available at: https://www.micromedexsolutions.com/ (cited: Nov/27/2025). 9. Lesko SM, Mitchell AA, Epstein M, et al. Heparin use as a risk factor for intraventricular hemorrhage in low-birth-weight infants. <i>N Engl J Med</i>. 1986; 314 (18): 1156 -60. 10. De Godoy J, De Marchi C, Silva M. Thrombosis of the Abdominal Aorta in a Newborn: Case Report and Review of Literature. <i>J Pediatr Surg</i>. 2003 Apr; 38(4):E11. 11. Nouri-Merchaoui S, Mahdhaoui N, Trabelsi S, et al. Spontaneous neonatal arterial thrombosis: a report of 4 neonates. <i>Arch Pediatr</i>. 2012 Apr; 19(4):413-8. 12. Moharir M, Shroff M, Stephens D, et al. Anticoagulants in pediatric cerebral sinovenous thrombosis: a safety and outcome study. <i>Ann Neurol</i>. 2010 May; 67(5):590-9. 13. Lau K, Stoffman J, Williams S, et al. Neonatal renal vein thrombosis: review of the English-language literature between 1992 and 2006. <i>Pediatrics</i>. 2007; 120(5):e1278-84. 14. Kosch A, Kuwertz-Broking E, Heller C, Kurnik K, Schobess R, et al. Renal venous thrombosis in neonates: prothrombotic risk factors and Long-term follow up. <i>Blood</i> 2004; 104:1356-60. 15. Morag I, Epelman M, Daneman A, et al. Portal vein thrombosis in the neonate: risk factors, course, and outcome. <i>J Pediatr</i>. 2006 Jun; 148(6):735-9. 16. Andrew M, Marzinotto V, Massicotte P, et al. Heparin therapy in pediatric patients: a prospective cohort study. <i>Pediatr Res</i>. 1994; 35 (1): 78 - 83. 17. Kulkarni M, Elsner C, Ouellet D, Zeldin R. Heparinised saline versus normal saline in maintaining patency of the radial artery catheter. <i>Can J Surg</i>. 1994 Feb; 37(1):37-42. 18. Butt W, Shan F, McDonnell G, Hudson I. Effect of heparin concentration and infusion rate on the patency of arterial catheters. <i>Crit Care Med</i>. 1987; 15 (3): 230 - 232.

19. Rais-Bahrami K, Karna P, Dolanski EA. Effect of fluids on life span of peripheral arterial lines. Am J Perinatol. 1990; 7 (2): 122 - 124.
20. Selldén H, Nilsson K, Larsson LE, Ekström-Jodal B. Radial arterial catheters in children and neonates: a prospective study . Crit Care Med. 1987; 15 (12): 1106 – 1109.
21. Andrew M, Marzinotto V, Massicotte P, Blanchette V, Ginsberg J, Brill-Edwards P, Burrows P, Benson L, Williams W, David M, Poon A. Heparin therapy in pediatric patients: a prospective cohort study. Pediatric Research. 1994 Jan;35(1):78-83.
22. Schmugge M, Risch L, Huber AR, Benn A, Fischer JE. Heparin-induced thrombocytopenia-associated thrombosis in pediatric intensive care patients. Pediatrics. 2002 Jan 1;109(1):e10.
23. Avila ML, Shah V, Brandao LR. Systematic review on heparin-induced thrombocytopenia in children: a call to action. Journal of Thrombosis and Haemostasis. 2013 Apr;11(4):660-9.
24. Pittiruti M, Bertoglio S, Scopettuolo G, Biffi R, Lamperti M, Dal Molin A, Panocchia N, Petrosillo N, Venditti M, Rigo C, DeLutio E. Evidence-based criteria for the choice and the clinical use of the most appropriate lock solutions for central venous catheters (excluding dialysis catheters): a GAVeCeLT consensus. The journal of vascular access. 2016 Nov;17(6):453-64.

Appendix

Table: Infusion rates when using heparin concentration **100 units/mL**

Rate (mL/hr)	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1
Weight (kg)	Approximate units/kg/hour									
0.5	20	40	60	80	100	120	140	160	180	200
1	10	20	30	40	50	60	70	80	90	100
1.5	7	13	20	27	33	40	47	53	60	67
2	5	10	15	20	25	30	35	40	45	50
2.5	4	8	12	16	20	24	28	32	36	40
3	3	7	10	13	17	20	23	27	30	33
3.5	3	6	9	11	14	17	20	23	26	29
4	3	5	8	10	13	15	18	20	23	25
4.5	2	4	7	9	11	13	16	18	20	22
5	2	4	6	8	10	12	14	16	18	20

$$\text{Rate (mL/hr)} = \frac{\text{Dose (unit/kg/hour)} \times \text{Weight (kg)}}{\text{Concentration (units/mL)}}$$

$$\text{Dose (unit/kg/hour)} = \frac{\text{Rate (mL/hr)} \times \text{Concentration (unit/mL)}}{\text{Weight (kg)}}$$

VERSION/NUMBER	DATE
Original 1.0	2/04/2026
Current 1.1	18/06/2026
REVIEW	2/04/2031

Authors Contribution

This standard concentration formulary has been developed by the ANMF standard concentration working group. The working group (in alphabetical order): Mohammad Irfan Azeem, Susannah Brew, Cindy Chen, Michelle Jenkins, Kerrie Knox, Rebecca O’Grady

Original author/s	Nilkant Phad, Srinivas Bolisetty, Juliana Teo, Bhavesh Mehta
-------------------	--

Evidence Review	Tim Schindler
Expert review	Juliana Teo
Nursing Review	Ben Emerson-Parker, Renae Gengaroli
Pharmacy Review	Susannah Brew, Kerrie Knox
ANMF Group contributors	Thao Tran, Michelle Jenkins, Rebecca Barzegar, Mohammad Irfan Azeem, Rebecca O’Grady, Cindy Chen, Celia Cunha da Silva, Kerrie Knox, Susannah Brew, Bryony Malloy, Renae Gengaroli, Samantha Hassall, Jutta van den Boom, Amber Seigel, Benjamin Emerson-Parker, Dianne Lee, Gloria Yoo, Simarjit Kaur, Emma Roylance, Trong Tran, Chales Tian
Final editing and review	Thao Tran, Srinivas Bolisetty
Facilitator	Srinivas Bolisetty

NEW RELEASE