Midazolam – Standard Concentration

Newborn use only

2026

<table>
<tr><td>Alert</td><td>S4 Drug – High risk medication causing significant patient harm when used in error.</td></tr>
<tr><td>Indication</td><td>Sedation
Seizures</td></tr>
<tr><td>Action</td><td>Intensify the physiological inhibitory mechanisms mediated by gamma-aminobutyric acid (GABA) by accumulation and occupation of benzodiazepine receptors. Anti-anxiety properties are related to increasing the glycine inhibitory neurotransmitter.</td></tr>
<tr><td>Drug type</td><td>Short acting benzodiazepine.</td></tr>
<tr><td>Trade name</td><td>Hypnovel, Midazolam Alphapharm, Midazolam Pfizer, Midazolam-Baxter, B. Braun Midazolam, Midazolam Accord, Midazolam Apotex.</td></tr>
<tr><td>Presentation</td><td>5mg/mL, 5mg/5mL, 50mg/10mL, and 15mg/3mL ampoules for IV and oral use.</td></tr>
<tr><td>Dose</td><td>
<table>
<tr><th>Method</th><th>Dose [1-5]</th></tr>
<tr><td>IV infusion for sedation</td><td>0.2–1 microgram/kg/minute</td></tr>
<tr><td>IV infusion for seizures</td><td>Loading dose: 150–200 microgram/kg over 3−5 minutes
Maintenance dose: 1–7 microgram/kg/minute</td></tr>
<tr><td>IV bolus</td><td>50 microgram/kg/dose every 2 hours when required.
(Dose range: 50–150 microgram/kg/dose)</td></tr>
<tr><td>IM injection</td><td>50 microgram/kg/dose every 4 hours when required.
(Dose range: 50–150 microgram/kg/dose)</td></tr>
<tr><td>Oral</td><td>250 microgram/kg as a single dose</td></tr>
<tr><td>Sublingual/buccal</td><td>200 microgram/kg as a single dose</td></tr>
<tr><td>Intranasal</td><td>200 microgram/kg per dose as a single dose
(Dose range: 200–300 microgram/kg/dose)</td></tr>
</table>
</td></tr>
<tr><td>Dose adjustment</td><td>Therapeutic hypothermia – No dose adjustment is required.[6]
ECMO – Increased volume of distribution but reduced renal clearance and accumulation of active metabolites over time. Higher dose may be required in early stages of ECMO. Close monitoring is recommended.[7]
Renal impairment – Limited data to recommend any dose adjustment.
Hepatic impairment – For repeated doses and IV infusion, reduction in dosage may be required.</td></tr>
<tr><td>Maximum dose</td><td></td></tr>
<tr><td>Total cumulative dose</td><td></td></tr>
<tr><td>Route</td><td>IV, IM, Oral, Sublingual.
Intranasal (not recommended due to nasal irritation; only under exceptional circumstances, e.g. acute refractory seizures with no alternate routes feasible).</td></tr>
<tr><td>Preparation</td><td>IV infusion
Note: Refer to Appendix for tables to assist with concentration selection.

Note: Use the smallest volume syringe available/suitable for drawing up the drug for the preparation. (e.g. for <1 mL draw up – use 1 mL syringe). For 10 mL syringe - Recommend using syringe that has markings at 0.2mL increments

The standard concentrations outlined in this monograph were developed taking into consideration a minimum patient weight of 0.4 kg and a minimum infusion rate of 0.1 mL/hour. For patients weighing less than 0.4 kg, further dilution may be required.

Prime the line as close to the entry point as possible to reduce the dead space – however, care should be taken not to deliver excess volume.

Weight suggestions for infusion concentrations below are a guide only. Clinicians may choose infusion concentration different to the suggested based on expected dose and the corresponding 24-hour fluid volumes</td></tr>
</table>

Midazolam concentration	50 microgram/mL	200 microgram/mL	400 microgram/mL	1000 microgram/mL
For seizures – Higher concentration (≥ 400 microgram/mL) is preferred for any infant weight.				
Infant weight (guide only)	<1 kg	1 to 3 kg	>3 kg	fluid restricted
0.2 microgram/kg/min is equal to	0.24 mL/kg/hour	0.06 mL/kg/hour	0.03 mL/kg/hour	0.012 mL/kg/hour
1 microgram/kg/min is equal to	1.2 mL/kg/hour	0.3 mL/kg/hour	0.15 mL/kg/hour	0.06 mL/kg/hour
50 microgram/kg bolus is equal to	1 mL/kg	0.25 mL/kg	0.125 mL/kg	0.05 mL/kg

Using Midazolam 5 mg/5mL strength (=1 mg/mL)

20mL Syringe

Draw up midazolam and add compatible fluid* as per table below to make a final volume of 20 mL

Midazolam concentration	50 microgram/mL	200 microgram/mL	400 microgram/mL	1000 microgram/mL
Volume of midazolam (1 mg/mL)	1 mL (=1 mg)	4 mL (=4 mg)	8 mL (=8 mg)	20 mL (=20 mg)
Volume of compatible fluid*	19 mL	16 mL	12 mL	Nil
Total volume	20 mL	20 mL	20 mL	20mL

* Compatible fluid: glucose 5%, glucose 10% or sodium chloride 0.9%

50mL Syringe

Draw up midazolam and add compatible fluid* as per table below to make a final volume of 50 mL

Midazolam concentration	50 microgram/mL	200 microgram/mL	400 microgram/mL	1000 microgram/mL
Volume of midazolam (1 mg/mL)	2.5 mL (=2.5 mg)	10 mL (=10 mg)	20 mL (=20 mg)	50 mL (=50 mg)
Volume of compatible fluid*	47.5 mL	40 mL	30 mL	Nil
Total volume	50 mL	50 mL	50 mL	50mL

* Compatible fluid: glucose 5%, glucose 10% or sodium chloride 0.9%

Using Midazolam 5 mg/mL strength

20mL Syringe

It is a 2-step dilution.

Step 1: Draw up midazolam and add compatible fluid* to make a diluted solution as per table below:

Midazolam concentration	50 microgram/mL	200 microgram/mL	400 microgram/mL	1000 microgram/mL
Volume of midazolam (5 mg/mL)	1 mL (=5 mg)		2 mL (=10 mg)	4 mL (=20 mg)
Volume of compatible fluid*	9 mL		8 mL	16 mL
Total volume	10 mL **(500 microgram/mL)**		10 mL **(1000 microgram/mL)**	20 mL **(1000 microgram/mL)**

Step 2: Draw up diluted midazolam and add compatible fluid* as per table below to make a final volume of 20 mL

Midazolam concentration	50 microgram/mL	200 microgram/mL	400 microgram/mL	1000 microgram/mL
Volume of diluted midazolam from step 1	2 mL (=1 mg)	8 mL (=4 mg)	8 mL (=8 mg)	20 mL (=20 mg)
Volume of compatible fluid*	18 mL	12 mL	12 mL	Nil
Total volume	20 mL	20 mL	20 mL	20mL

* Compatible fluid: glucose 5%, glucose 10% or sodium chloride 0.9%

50mL Syringe

It is a 2-step dilution.

Step 1: Draw up midazolam and add compatible fluid* to make a diluted solution as per table below:

Midazolam concentration	50 microgram/mL	200 microgram/mL	400 microgram/mL	1000 microgram/mL
Volume of midazolam (5 mg/mL)	1 mL (=5 mg)	2 mL (=10 mg)	4 mL (=20 mg)	10 mL (=50 mg)
Volume of compatible fluid*	9 mL	Nil	Nil	Nil
Total volume	10 mL **(500 microgram/mL)**	2 mL **(5mg/mL)**	4 mL **(5mg/mL)**	10 mL **(5mg/mL)**

Step 2: Draw up midazolam and add compatible fluid* as per table below to make a final volume of 50 mL

Midazolam concentration	50 microgram/mL	200 microgram/mL	400 microgram/mL	1000 microgram/mL
Volume of diluted midazolam from step 1	5 mL (=2.5 mg)	2 mL (=10 mg)	4 mL (=20 mg)	10 mL (=50 mg)
Volume of compatible fluid*	45 mL	48 mL	46 mL	40 mL
Total volume	50 mL	50 mL	50 mL	50mL

* Compatible fluid: glucose 5%, glucose 10% or sodium chloride 0.9%

IV bolus, IM, oral, sublingual

Using 5 mg/5mL strength (= 1 mg/mL)

Draw up 1 mL (1000 microgram of midazolam) and add 4 mL of sodium chloride 0.9% to make final volume of 5 mL with a concentration of 200 microgram/mL.

Using 5 mg/mL or 15 mg/3mL strength (= 5 mg/mL)

Draw up 0.4 mL (2000 microgram of midazolam) and add 9.6 mL of sodium chloride 0.9% to make final volume of 10 mL with a concentration of 200 microgram/mL.

Intranasal

Using the 5 mg/5mL strength (= 1 mg/mL)

Consider using undiluted midazolam

Using 5 mg/mL or 15 mg/3mL strength (= 5 mg/mL)

Infant <3 kg

Draw up 0.2 mL (1000 microgram of midazolam) and add 0.8 mL of sodium chloride 0.9% to make a final volume of 1 ml and concentration of 1 mg/mL (1000 microgram/mL)
Recommended maximum volume in each nostril: 0.3 mL. Larger volumes may end up in the nasopharynx.[24]

	Infant ≥3 kg Draw up 0.4 mL (2000 microgram of midazolam) and add 0.6 mL of sodium chloride 0.9% to make a final volume of 1 ml and concentration of 2 mg/mL (=2000 microgram/mL). Recommended maximum volume in each nostril: 0.3 mL. Larger volumes may end up in the nasopharynx.[24]
Administration	**IV infusion:** continuous infusion via a syringe pump. Change solution every 24 hours. **IV bolus:** slow push over 10 minutes.[8] **Oral, sublingual:** Only solutions prepared from **plastic** IV ampoules may be used for oral or sublingual administration. **IM:** Inject deep into a large muscle. **Intranasal:** **Direct administration:** Drop dose into alternating nostrils over 15 seconds. Absorption is rapid; maximum effect in 10 minutes and duration up to 2 hours. May be irritating to nasal mucosa. **Mucosal atomisation device (MAD):** • Attach MAD to the end of a 1 mL Luer- lock syringe and prime the device with the midazolam solution to the prescribed dose. • Insert the MAD loosely into the nostril to form a seal, preventing expulsion of fluid. • Briskly compress the syringe plunger to allow for maximal coverage of nasal mucosa with atomised particles.
Monitoring	Apnoea, respiratory depression. Blood pressure. Level of sedation.
Contraindications	Known hypersensitivity to midazolam.
Precautions	In preterm infants, especially in extreme preterm, midazolam half-life is increased from 4−6 hours in term neonates up to 22 hours in premature infants. It is longer with impaired liver function. Caution when concurrently used with opioids − midazolam interacts with other central nervous system (CNS) depressants and may increase the risk of drowsiness, respiratory depression, and hypotension. Withdraw slowly after chronic administration as abrupt discontinuation may precipitate withdrawal seizures. Caution in neonates with renal and hepatic impairment − increased sensitivity to CNS effects; use doses at lower end of the range. Rapid IV infusion may result in hypotension, respiratory depression, or seizure.
Drug interactions	Concurrent administration with erythromycin promotes accumulation. Xanthines may decrease the anaesthetic/sedative effect of benzodiazepines. Care needs to be taken with adding or withdrawing caffeine or aminophylline.
Adverse reactions	Hypotension and reduced cardiac output, particularly when used in combination with fentanyl. Respiratory depression and apnoea. Hypersalivation. Nasal discomfort (with intranasal route). Seizure-like myoclonus (more common in premature neonates receiving via intravenous route).[9]
Overdose	Flumazenil is a specific benzodiazepine receptor antagonist. Dose: 10 microgram/kg IV bolus followed by repeat boluses (1-2 min) or 5 microgram/kg/min IV infusion until sedation reversed or maximum dose of 50 microgram/kg is reached.[10]
Compatibility	Fluids: Glucose 5%, glucose 10%, sodium chloride 0.9%, and sodium chloride 0.45%. Y-site[11]: Amino acid solutions. Acetaminophen, amikacin, amiodarone, atracurium, atropine, aztreonam, calcium chloride, calcium gluconate, caspofungin, cefazolin, cefotaxime, cefoxitin, ceftriaxone, ciprofloxacin, dexmedetomidine, digoxin, diltiazem, dopamine, doxycycline, enalaprilat, epinephrine, erythromycin lactobionate, fentanyl, fluconazole, folic acid (as sodium salt), gentamicin, glycopyrrolate, heparin, isoproterenol, ketamine, labetolol, lidocaine, linezolid, lorazepam, magnesium sulfate, metronidazole, milrinone, morphine hydrochloride, morphine sulfate, multiple vitamin injection, naloxone, nitroglycerin, nitroprusside sodium, norepinephrine, octreotide, oxacillin, pamidronate, pancuronium, papaverine, penicillin G potassium, penicillin G sodium, pentoxyfylline, piperacillin, potassium chloride, procainamide, propranolol, protamine sulfate, pyridoxine, ranitidine, remifentanil,

	rocuronium, streptokinase, theophylline, ticarcillin, ticarcillin-clavulanate, tobramycin, urokinase, vancomycin, vasopressin, vecuronium, and verapamil. **Variable compatibility[11]:** amoxicillin-clavulanate, clindamycin, clonidine, dobutamine, furosemide, hydralazine, imipenem-cilastatin, insulin, regular, methylprednisolone sodium succinate, pantoprazole, propofol, SMOFlipid (up to 0.5 mg/mL midazolam concentration)[23], and sodium acetate.
Incompatibility	Fluids: No information. Y-site: Aciclovir, albumin, aminophylline, amoxicillin, amphotericin B cholesteryl sulfate complex, amphotericin B conventional colloidal, amphotericin B lipid complex, amphotericin B liposome, ampicillin, atenolol, azathioprine, azithromycin, cefepime, ceftazidime, chloramphenicol, cloxacillin, dexamethasone, diazepam, diazoxide, epoetin alfa, esomeprazole, flucloxacillin, fluorouracil, ganciclovir, hydrocortisone sodium succinate, ibuprofen lysine, indomethacin, omeprazole, phenobarbital (phenobarbitone), phenytoin, piperacillin-tazobactam, potassium acetate, sodium bicarbonate, sulfamethoxazole-trimethoprim, and thiopental.[11]
Stability	Diluted solution: Store at 2−8°C and use within 24 hours.
Storage	Midazolam Apotex, Midazolam-Baxter: Store below 30°C. Protect from light. B. Braun Midazolam, Hypnovel, Midazolam Alphapharm: Store below 25°C. Protect from light. Midazolam Pfizer: Store below 25°C. Protect from light. Unopened ampoules will be suitable for use for up to 8 months after the foil sachet has been opened, if protected from light. Schedule 4D (S4D) medication. Store in dangerous drug safe and record use in S4D register.
Excipients	Sodium chloride, hydrochloric acid, sodium hydroxide, and water for injections.
Special comments	Flumazenil is a specific benzodiazepine antagonist and may be used (very limited experience in the neonate) to rapidly reverse respiratory depression − 10 microgram/kg/dose IV push. May repeat every minute for up to 4 more doses.
Evidence	**Efficacy** **Sedation** There is insufficient evidence to promote the routine use of intravenous midazolam infusion as a sedative for neonates undergoing intensive care. Although all studies included in the review reported better sedation, none of the scales used had been validated in preterm infants and thus the effectiveness could not be evaluated.[12] (Level 1, Grade B). Intranasal midazolam for sedation: In a randomised control trial Milesi et al administered intranasal midazolam to 27 neonates of mean gestational age 27 weeks in the delivery room prior to intubation. The neonates allocated to the nasal midazolam arm received 0.1 mg/kg (0.1 mL/kg) of midazolam in each nostril. Nasal midazolam was more efficient than nasal Ketamine (89% vs 58%; p<0.01) for sedation. The haemodynamic and respiratory effects of both drugs were comparable.[4] Ku et al described a retrospective cohort of 18 infants receiving 20 intranasal doses of midazolam.[13] The median gestational age of infants at birth was 27 weeks and postnatal age was 34 days. The median dose was 0.1 mg/kg (0.1 -0.2). All the infants tolerated the medication well and none developed hypotension, bradycardia, or died.[14] **Seizures** In a small (n=39) retrospective cohort study, Conde et al reported high effectiveness of intravenous midazolam as the first line anti-seizure medication to control electrographic seizures secondary to HIE, perinatal stroke, or idiopathic in term neonates. Midazolam was administered as a 150 microgram/kg IV bolus followed by a continuous infusion 1-18 microgram/kg/min. The therapeutic response was measured as total seizure burden, maximum ictal fraction (MIF), and EEG background. Seizure activity was considered controlled if within 4 hours of reaching the maximum dose of midazolam the MIF was less than 3 minutes with > 80% reduction in average ictal fraction and there was no recurrence of EEG seizures. In 61% neonates, adequate seizure control was achieved with midazolam alone and in a further 38% neonates in combination with lidocaine. Abnormal neurodevelopment (n=10) beyond 18 months correlated with the EEG background at the beginning and 24-48 hours after seizure control but not with therapeutic response.[3] Midazolam was effective in neonates with refractory seizures that did not respond to phenytoin or pentobarbital (pentobarbitone).[2] (Level IV, Grade D).

	Intranasal midazolam for seizures: In a randomised study, Fisgin et al administered 0.2 mg/kg midazolam intranasally to 16 participants aged 0-24 months over 30 seconds using an injector. The age of youngest participants was 1 month but the number of participants of age 1 month was not clear. In 87% of the participants in the nasal midazolam group the seizures were terminated compared to 60% in the rectal diazepam group. Authors reported no major adverse events following intranasal midazolam.[5,15] **Safety** There is limited evidence of the risk/benefit ratio for sedative use in mechanically ventilated preterm neonates. The EPIPAGE-2 cohort study reported better survival and a comparable 2-year development profile in preterm infants who received opioid and/or midazolam infusion during initial mechanical ventilation during the first week after birth. This study included infants born between completed 23- and 31-week gestation who were intubated within 1 hour of birth and remained intubated > 24 hours. The treatment group infants (n=239) received a continuous infusion of opioids and/or midazolam during this first intubation period or day 7. The control group infants either did not receive midazolam during their stay in the NICU (n= 290) or received it after their first extubation or day 7 (n=113). The cumulative mean duration of midazolam infusion was 4 (IQR 2-11) days. Neurodevelopment was assessed in 94 infants in the treatment arm and 107 infants in the control arm.[16] One study showed a statistically significant higher incidence of adverse neurological events (death, grade III or IV IVH, and PVL) and meta-analysis of data from two studies showed a statistically significant longer duration of NICU stay in the midazolam group compared to the placebo group.[12] (Level1, Grade B). Administration of midazolam in ventilated premature infants causes significant changes in cerebral oxygenation and hemodynamics, which might be harmful.[17] (Level III, Grade C). Intravenous bolus doses of midazolam in association with fentanyl should be used with great caution in the newborn, especially if very premature or with unstable blood pressure.[18] (Level IV, Grade D). Sedation with midazolam has a transient effect on the background aEEG activity.[19] (Level III, Grade C). Flumazenil is a specific benzodiazepine receptor antagonist. In one small, randomised control trial 40 healthy 3–12-year-old children, 500 microgram/kg oral midazolam was used for premedication and additional IV 500 microgram/kg for induction of anesthesia for circumcision surgery in addition to nitrous oxide, halothane, and bupivacaine. Reversal of sedation was attempted using a blinded study medication 3 min after child returned to recovery after surgery. Children who received flumazenil woke up 4 times faster compared to the placebo group. The average total dose of flumazenil administered was 24(±19) microgram/kg.[10] **Pharmacokinetics** Midazolam is highly protein bound with an elimination half-life of 4–6 hours in term neonates and a variable half-life (up to 22 hours) in premature neonates and those with impaired hepatic function. Bioavailability is approximately 36% with oral administration and 50% with sublingual and intranasal administration.[20] (Level III, Grade C). Pharmacokinetic data favours low dose of IV infusion for sedation in very preterm neonates compared to more mature neonates.[1]
Practice points	
References	1. Völler S, Flint RB, Beggah F, et. Recently Registered Midazolam Doses for Preterm Neonates Do Not Lead to Equal Exposure: A Population Pharmacokinetic Model. J Clin Pharmacol. 2019 Oct;59(10):1300-1308. 2. Castro Conde JR, Hernandez Borges AA, Domenech Martinez E, et al. Midazolam in neonatal seizures with no response to phenobarbital. Neurology. Mar 8, 2005;64(5):876–879. 3. Castro Conde JR, González Campo C, et al. High Effectiveness of Midazolam and Lidocaine in the Treatment of Acute Neonatal Seizures. J Clin Neurophysiol. 2024 Jul 1;41(5):450-457. 4. Milési C, Baleine J, Mura T, et al. Nasal midazolam vs ketamine for neonatal intubation in the delivery room: a randomised trial Arch Dis Child Fetal Neonatal Ed 2018; 103: F221–F226. 5. Fişgin T, Gürer Y, Senbil N, et al. Nasal midazolam effects on childhood acute seizures. J Child Neurol. 2000 Dec; 15(12):833-5. 6. Favié, Laurent M A et al. "Phenobarbital, Midazolam Pharmacokinetics, Effectiveness, and Drug-Drug Interaction in Asphyxiated Neonates Undergoing Therapeutic Hypothermia." Neonatology vol. 116, 2 (2019): 154-162.

7. Raffaeli G, Pokorna P, Allegaert K, et al. Drug Disposition and Pharmacotherapy in Neonatal ECMO: From Fragmented Data to Integrated Knowledge. Front Pediatr. 2019; 7:360. Published 2019 Sep 3. doi:10.3389/fped.2019.00360
8. Van Den Broek MP, Van Straaten HL, Huitema AD, et al. Anticonvulsant effectiveness and hemodynamic safety of midazolam in full-term infants treated with hypothermia. Neonatology. 2015 Jan 8;107(2):150-6.
9. Gupta MK, Mondkar JA, Hegde D. Paradoxical Reaction to Midazolam in Preterm Neonates: A Case Series. Indian J Crit Care Med. 2018 Apr;22(4):300-302.
10. Jones RD, Lawson AD, Andrew LJ, et al. Antagonism of the hypnotic effect of midazolam in children: a randomized, double-blind study of placebo and flumazenil administered after midazolam-induced anaesthesia. Br J Anaesth. 1991 Jun;66(6):660-6.
11. Micromedex online. Midazolam. Accessed on 21 April 2021.
12. Ng E, Taddio A, Ohlsson A. Intravenous midazolam infusion for sedation of infants in the neonatal intensive care unit. Cochrane Database of Systematic Reviews 2017, Issue 1. Art. No.: CD002052.
13. Ku LC, Simmons C, Smith PB, et al. Intranasal midazolam and fentanyl for procedural sedation and analgesia in infants in the neonatal intensive care unit. J Neonatal Perinatal Med. 2019;12(2):143-148.
14. Lawrence C. Ku, Catherine Simmons, Brian Smith, et al. Pediatrics Jan 2018, 141 (1 Meeting Abstract) 532; DOI: 10.1542/peds.141.1_MeetingAbstract.532.
15. Humphries LK, Eiland LS. Treatment of acute seizures: is intranasal midazolam a viable option? J Pediatr Pharmacol Ther. 2013 Apr;18(2):79-87.
16. de Tristan MA, Martin-Marchand L, Roué JM, et al. Association of Continuous Opioids and/or Midazolam During Early Mechanical Ventilation with Survival and Sensorimotor Outcomes at Age 2 Years in Premature Infants: Results from the French Prospective National EPIPAGE 2 Cohort. J Pediatr. 2021 May; 232:38-47.
17. Van Alfen-van der Velden AA, Hopman JC, Klaessens JH, Feuth T, Sengers RC, Liem KD. Effects of midazolam and morphine on cerebral oxygenation and hemodynamics in ventilated premature infants. Biology of the Neonate. 2006;90(3):197–202.
18. Burtin P, Daoud P, Jacqz-Aigrain E, Mussat P, Moriette G. Hypotension with midazolam and fentanyl in the newborn. Lancet. Jun 22, 1991;337(8756):1545–1546
19. Bernet V, Latal B, Natalucci G, Doell C, Ziegler A, Wohlrab G. Effect of sedation and analgesia on postoperative amplitude-integrated EEG in newborn cardiac patients. Pediatr Res. Jun 2010;67(6):650–655.
20. De Wildt SN, Kearns GL, Hop WC, Murry DJ, Abdel-Rahman SM, van den Anker JN. Pharmacokinetics and metabolism of intravenous midazolam in preterm infants. Clin Pharmacol Ther. 2001 Dec;70(6):525–31.
21. Taketomo CK, Hodding JH, Kraus DM, American Pharmacists Association. Pediatric and neonatal dosage handbook. Hudson, Ohio: Lexi-Comp: American Pharmacists Association; 2015.
22. MerativeTM Micromedex® Complete IV Compatibility (electronic version). Merative, Ann Arbor, Michigan, USA. Available at: https://www.micromedexsolutions.com/ (cited: Aug/15/2024).
23. Senarathna SMDKG, Strunk T, Petrovski M, Woodland S, Martinez J, Chuang VTG, Batty KT. Physical compatibility of lipid emulsions and intravenous medications used in neonatal intensive care settings. Eur J Hosp Pharm 2023; In press; doi:10.1136/ejhpharm-2023-003870.
24. Pansini V, Curatola A, Gatto A, Lazzareschi I, Ruggiero A, Chiaretti A. Intranasal drugs for analgesia and sedation in children admitted to pediatric emergency department: a narrative review. Annals of translational medicine. 2021 Jan;9(2).

Appendix

Infusion tables to assist concentration selection

Table 1: Infusion rates when using midazolam concentration **50 microgram/mL**

Rate (mL/hr)	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1
Weight (kg)	**Approximate microgram/kg/minute**									
0.5	0.17	0.33	0.5	0.67	0.83	1	1.2	1.3	1.5	1.7
1	0.08	0.17	0.25	0.33	0.42	0.5	0.58	0.67	0.75	0.83
1.5	0.06	0.11	0.17	0.22	0.28	0.33	0.39	0.44	0.5	0.56
2	0.04	0.08	0.13	0.17	0.21	0.25	0.29	0.33	0.38	0.42
2.5	0.03	0.07	0.1	0.13	0.17	0.2	0.23	0.27	0.3	0.33
3	0.03	0.06	0.08	0.11	0.14	0.17	0.19	0.22	0.25	0.28
3.5	0.02	0.05	0.07	0.1	0.12	0.14	0.17	0.19	0.21	0.24
4	0.02	0.04	0.06	0.08	0.1	0.13	0.15	0.17	0.19	0.21
4.5	0.02	0.04	0.06	0.07	0.09	0.11	0.13	0.15	0.17	0.19
5	0.02	0.03	0.05	0.07	0.08	0.1	0.12	0.13	0.15	0.17

Table 2: Infusion rates when using midazolam concentration **200 microgram/mL**

Rate (mL/hr)	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1
Weight (kg)	**Approximate micrograms/kg/minute**									
0.5	0.67	1.33	2	2.7	3.3	4	4.7	5.3	6	6.7
1	0.33	0.67	1	1.3	1.7	2	2.3	2.7	3	3.3
1.5	0.22	0.44	0.67	0.89	1.1	1.3	1.6	1.8	2	2.2
2	0.17	0.33	0.5	0.67	0.83	1	1.2	1.3	1.5	1.7
2.5	0.13	0.27	0.4	0.53	0.67	0.8	0.93	1.1	1.2	1.3
3	0.11	0.22	0.33	0.44	0.56	0.67	0.78	0.89	1	1.1
3.5	0.1	0.19	0.29	0.38	0.48	0.57	0.67	0.76	0.86	0.95
4	0.08	0.17	0.25	0.33	0.42	0.5	0.58	0.67	0.75	0.83
4.5	0.07	0.15	0.22	0.3	0.37	0.44	0.52	0.59	0.67	0.74
5	0.07	0.13	0.2	0.27	0.33	0.4	0.47	0.53	0.6	0.67

Table 3: Infusion rates when using midazolam concentration **400 microgram/mL**
Doses in black colour are within sedation range, doses in orange are within seizure range.

Rate (mL/hr)	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1
Weight (kg)	**Approximate micrograms/kg/minute**									
0.5	1.3	2.7	4	5.3	6.7	8	9.3	11	12	13
1	0.67	1.3	2	2.7	3.3	4	4.7	5.3	6	6.7
1.5	0.44	0.89	1.3	1.8	2.2	2.7	3.1	3.6	4	4.4
2	0.33	0.67	1	1.3	1.7	2	2.3	2.7	3	3.3
2.5	0.27	0.53	0.8	1.1	1.3	1.6	1.9	2.1	2.4	2.7
3	0.22	0.44	0.67	0.89	1.1	1.3	1.6	1.8	2	2.2
3.5	0.19	0.38	0.57	0.76	0.95	1.14	1.33	1.5	1.7	1.9
4	0.17	0.33	0.5	0.67	0.83	1	1.17	1.3	1.5	1.7
4.5	0.15	0.3	0.44	0.59	0.74	0.89	1	1.2	1.3	1.5
5	0.13	0.27	0.4	0.53	0.67	0.8	0.93	1.1	1.2	1.3

Table 4: Infusion rates when using midazolam concentration **1000 microgram/mL (1 mg/mL**)
Doses in black colour are within sedation range, doses in orange are within seizure range.

Rate (mL/hr)	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1
Weight (kg)	**Approximate micrograms/kg/minute**									
0.5	3.3	6.7	10	13	17	20	23	27	30	33
1	1.7	3.3	5	6.7	8.3	10	12	13	15	17
1.5	1.1	2.2	3.3	4.4	5.6	6.7	7.8	8.9	10	11
2	0.83	1.7	2.5	3.3	4.2	5	5.8	6.7	7.5	8.3
2.5	0.67	1.3	2	2.7	3.3	4	4.7	5.3	6	6.7
3	0.56	1.1	1.7	2.2	2.8	3.3	3.9	4.4	5	5.6
3.5	0.48	0.95	1.4	1.9	2.4	2.9	3.3	3.8	4.3	4.8
4	0.42	0.83	1.3	1.7	2.1	2.5	2.9	3.3	3.8	4.2
4.5	0.37	0.74	1.1	1.5	1.9	2.2	2.6	3	3.3	3.7
5	0.33	0.67	1	1.3	1.7	2	2.3	2.7	3	3.3

$$\text{Rate (mL/hr)} = \frac{60 \times \text{Dose (microgram/kg/min)} \times \text{Weight (kg)}}{\text{Concentration (microgram/mL)}}$$

$$\text{Dose (microgram/kg/min)} = \frac{\text{Rate (mL/hr)} \times \text{Concentration (microgram/mL)}}{\text{Weight (kg)} \times 60}$$

VERSION/NUMBER	**DATE**
Original 1.0	28/10/2025
Current 1.1	08/10/2026
REVIEW	08/10/2031

This standard concentration formulary has been developed by the ANMF standard concentration working group.
The working group (in alphabetical order): Celia Cunha da Silva, Cindy Chen, Dianne Lee, Gloria Yoo, Kerrie Knox, Michelle Jenkins, Mohammad Irfan Azeem, Rebecca O'Grady, Simarjit Kaur and Susanah Brew.

Authors Contribution of the current version

Current version authors	Nilkant Phad, Srinivas Bolisetty
Evidence Review	Nilkant Phad, Srinivas Bolisetty
Expert review	
Nursing Review	Renae Gengaroli, Celia Cunha da Silva
Pharmacy Review	Kerrie Knox, Susannah Brew, Mohammad Irfan Azeem
ANMF Group contributors	Nilkant Phad, Bhavesh Mehta, Amber Seigel, Rebecca Barzegar, Jutta van den Boom, Mohammed Irfan Azeem, Michelle Jenkins, Kerrie Knox, Cindy Chen, Benjamin Emerson-Parker, Renae Gengaroli, Dianne Lee, Gloria Yoo, Bryony Malloy, Celia Cunha da Silva, Simarjit Kaur, Thao Tran
Final editing	Srinivas Bolisetty, Celia Cunha da Silva
Electronic version	Thao Tran, Helen Huynh, Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty